

NOAA OAR Special Report

PMEL Tsunami Forecast Series: Vol. NNNN
**Development of a Tsunami Forecast Model for Arena
Cove, California**

Michael C. Spillane ^{1,2}

¹Joint Institute for the Study of the Atmosphere and Ocean (JISAO),
University of Washington, Seattle, WA

²NOAA/Pacific Marine Environmental Laboratory (PMEL), Seattle, WA

April 2011

NOTICE from NOAA

Mention of a commercial company or product does not constitute an endorsement by NOAA/OAR. Use of information from this publication concerning proprietary products or the tests of such products for publicity or advertising purposes is not authorized. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Oceanic and Atmospheric Administration.

Contribution No. XXXX from NOAA/Pacific Marine Environmental Laboratory

Also available from the National Technical Information Service (NTIS)
(<http://www.ntis.gov>)

Contents

List of Figures

List of Tables

Foreword

Abstract

1.0 Background and Objectives

- 1.1 The Setting
- 1.2 Natural Hazards
- 1.3 Tsunami Warning and Risk Assessment

2.0 Forecast Methodology

- 2.1 The Tsunami Model
- 2.2 The SIFT Forecast System

3.0 Model Development

- 3.1 Digital Elevation Models
- 3.2 Tides and Sea Level Variation
- 3.3 The CFL Condition and other considerations for grid design
- 3.4 Specifics of the model grids
- 3.5 Model Run Input and Output Files

4.0 Results and Discussion

- 4.1 The “Null” Tests
- 4.2 The Extreme Case Tests
- 4.3 Model Validation with Historical Events
- 4.4 Further Historical Simulations
- 4.5 The Mendocino Earthquake of April 25, 1992
- 4.6 The Honshu Tsunami of March 11, 2011
- 4.7 Simulation of further Extreme Events

5.0 Conclusions

6.0 Acknowledgements

7.0 References

FIGURES

Appendix A

- A1. Reference Model Input (*.in) File for Arena Cove, CA
- A2. Forecast Model Input (*.in) File for Arena Cove, CA

Appendix B Propagation Database: Pacific Ocean Unit Sources

Appendix C Synthetic Testing Report

Glossary

DRAFT

List of Figures

- Figure 1.** The Point Arena area of southern Mendocino County, CA.
- Figure 2.** Views of present-day Arena Cove and its appearance in the early 1900's.
- Figure 3.** Extract from the oblique 3-D view of the Arena Cove DEM, provided by NGDC; sites of potential inundation identified by CalEMA are highlighted in red.
- Figure 4.** Distribution of the historical tsunami sources employed for the development of the Arena Cove forecast model. Those highlighted in red are more extensively investigated using the reference model.
- Figure 5.** A sample interval from the Arena Cove tsunami-capable tide gage.
- Figure 6.** The setting of the Arena Cove model.
- Figure 7.** Nested grid representation for the Reference Model (RM). Rectangles are drawn, when appropriate, to indicate the placement of the next nesting level.
- Figure 8.** Nested grid representation for the Forecast Model (FM). Rectangles are drawn, when appropriate, to indicate the placement of the next nesting level.
- Figure 9.** Illustration of the appearance of model instability (upper panel.) The RM C-grid bathymetry in this case required minor modifications, following which the “null” test gave satisfactory results.
- Figure 10.** Locations of synthetic tsunami scenarios employed in model development.
- Figure 11.** Comparison of Reference (RM) and Forecast (FM) model results for the ACSZ 56-65 synthetic mega-event, representing the Cascadia Subduction Zone. Time series at the tide gage location in Arena Cove are shown in the upper panel. The lower panels contrast the RM and FM amplitude and velocity fields at the time indicated. Insets are enlargements of Arena Cove.
- Figure 12.** As in Figure 11, but for the KISZ 01-10 scenario representing Kamchatka.
- Figure 13.** As in Figure 11, but for the NTSZ 30-30 scenario representing Samoa.
- Figure 14.** Comparison of the Reference and Forecast Model response to a synthetic moderate event at NTSZ B36 near Samoa.
- Figure 15.** Comparison of the RM and FM response for the historical Unimak-1946 tsunami (prior to Arena Cove tide gage.).
- Figure 16.** Comparison of the RM and FM response for the historical Alaska-1964 tsunamis (prior to Arena Cove tide gage.)
- Figure 17.** Comparison of RM and FM hind casts of the Kuril 2006 event to sea level fluctuations in Arena Cove, observed by the tsunami-capable tide gage.
- Figure 18.** Comparison of RM and FM hind casts of the Samoa 2009 event to sea level fluctuations in Arena Cove.
- Figure 19.** Comparison of RM and FM hind casts of the Chile 2010 event to sea level fluctuations in Arena Cove.
- Figure 20.** Simulated model response (FM) to the historical Sanriku 1896, Kamchatka 1952, Andreanof 1957 and Chile 1960 events. Proxy observations (in red) are provided where available from other California locations. The Sanriku 1896 result is not expected to match the observed run-up at nearby Mendocino. The state of the tide is shown and times are given as UTC.
- Figure 21.** Model response (FM) to the East Kuril 1994, Chile 1995, Kuril 1995 and Irian Jaya 1996 events. Tide gage data (in red) where available is from Arena Cove.

- Figure 22.** Model response (FM) to the Andreanof 1996, Peru 2001, Hokkaido 2003, and Rat Island 2003 events. Tide gage data (in red) are from Arena Cove.
- Figure 23.** Model response (FM) to the Tonga 2006, Kuril 2007, Solomons 2007, and Peru 2007 events. Tide gage data (in red) are from Arena Cove.
- Figure 24.** Model response (FM) to the Chile 2007, Papua-New Guinea 2009, Solomons 2009, and Vanuatu 2009 events. Tide gage data (in red) are from Arena Cove.
- Figure 25.** Model response (FM) to the Samoa 2009 and Chile 2010 events. Tide gage data (in red) are from Arena Cove.
- Figure 26.** Seismicity in the vicinity of Cape Mendocino and the source mechanisms of notable recent earthquakes, adapted from USGC/NEIC products. The lower panel shows the poor agreement between the model and Arena Cove tide gage observations that is likely due to inadequate source representation in the model.
- Figure 27.** Comparison of FM real-time forecast and RM hindcast of the Honshu 2011 event with sea level observations in Arena Cove.
- Figure 28.** Predicted maximum sea level (from the FM) at the Arena Cove tide gage for the “mega-event” scenarios described in Table 6. Great circle routes are shown in red (with distances in km), and black arrows indicate the normal to the strike direction.
- Figure 29.** Comparison of RM and FM predictions for inundation of the Arena Cove / Manchester Beach region for selected mega-tsunami scenarios, and (lower right) for the ensemble employed in the CalEMA study.

List of Tables

- Table 1a.** Source characterization for historical tsunami events employed in Arena Cove model testing. Those in bold text were used in RM/FM inter-comparison. Sources identified as “ad hoc” may not be identically defined in other Forecast Model reports.
- Table 1b.** Supplementary historical tsunami events employed for Arena Cove forecast model testing. Those identified as “ad hoc” and may not be identically implemented in other Forecast Model reports.
- Table 2.** The main features of the Arena Cove Digital Elevation Model.
- Table 3.** Tidal characteristics of the Arena Cove, CA Tide Gage (9416841).
- Table 4.** Specifics of the grids and model parameters employed to model Arena Cove, CA. “EWxNS” denotes the number of grid values in the zonal (East to West) and meridional (North to South) directions respectively.
- Table 5.** Grid file names and grid-related parameters. The time steps for the A and B-grids must be integer multiples of the basic time step chosen for the C-grid.
- Table 6.** Description of the synthetic tsunami scenarios. Details of the source zones and specifics of the individual unit sources are provided in Appendix B.

This page is intentionally blank.

DRAFT

PMEL Tsunami Forecast Series: Vol. NNNN
**Development of a Tsunami Forecast Model for Arena
Cove, California**

Michael C. Spillane^{1,2}

Foreword

Tsunamis have been recognized as a potential hazard to United States coastal communities since the mid-twentieth century, when multiple destructive tsunamis caused damage to the states of Hawaii, Alaska, California, Oregon, and Washington. In response to these events, the United States, under the auspices of the National Oceanic and Atmospheric Administration (NOAA), established the Pacific and Alaska Tsunami Warning Centers, dedicated to protecting United States interests from the threat posed by tsunamis. NOAA also created a tsunami research program at the Pacific Marine Environmental Laboratory (PMEL) to develop improved warning products.

The scale of destruction and unprecedented loss of life following the December 2004 Sumatra tsunami served as the catalyst to refocus efforts in the United States on reducing tsunami vulnerability of coastal communities, and on 20 December 2006, the United States Congress passed the “Tsunami Warning and Education Act” under which education and warning activities were thereafter specified and mandated. A “tsunami forecasting capability based on models and measurements, including tsunami inundation models and maps.” is a central component for the protection of United States coastlines from the threat posed by tsunamis. The forecasting capability for each community described in the PMEL Tsunami Forecast Series is the result of collaboration between the National Oceanic and Atmospheric Administration office of Oceanic and Atmospheric Research, National Weather Service, National Ocean Service, National Environmental Satellite Data and Information Service, the University of Washington’s Joint Institute for the Study of the Atmosphere and Ocean, National Science Foundation, and United States Geological Survey.

Abstract. Operational tsunami forecasting by NOAA’s Tsunami Warning Centers relies on the detection of tsunami wave trains in the open ocean, the inversion of these data (telemetered via satellite) to quantify their source characteristics, and real-time modeling of the impact on threatened coastal communities. The latter phase of the process involves, for each such community, a pre-tested Forecast Model capable of predicting the impact, in terms of inundation and dangerous inshore currents, with sufficient resolution and within the time constraints appropriate to an emergency response.

In order to achieve this goal, considerable advance effort is required to tune each forecast model to the specific bathymetry and topography, both natural and manmade, of the impact area, and to validate its performance with a broad set of tsunami sources. Where possible the validation runs should replicate observed responses to historical events, but the sparse instrumental record of these rare but occasionally devastating occurrences

dictates that comprehensive testing should include a suite of scenarios that represent potential future events.

During the forecast model design phase, and in research mode outside the pressures of an emergency situation, more detailed and slower-running models can be investigated. Such a model, referred to as a Reference Model, represents the most credible numerical representation of tsunami response for the study region, using the most detailed bathymetry available and without the run-time constraint of operational use. Once a reference model has been developed, the process of forecast model design is to determine where efficiencies can be gained, through reducing the grid resolution and increasing the model time step, while still adequately representing the salient features of the full solution.

This report documents the reference and forecast model development for Arena Cove, a small inlet south of the rocky Arena Point headland in southern Mendocino County, CA. Both the cove and the headland serve as reference points in coastal reports and are tourist venues but, while several tsunami have been detected by the tide gage there, no injury or infrastructure damage have been reported to date. The Manchester Beach area, north of Point Arena, is low-lying and subject to inundation and has been included in the model domain.

¹ Joint Institute for the Study of the Atmosphere and Ocean (JISAO), University of Washington, Seattle, WA

²NOAA/Pacific Marine Environmental Laboratory (PMEL), Seattle, WA

1.0 Background and Objectives

1.1. The Setting

Arena Cove is a semi-circular indentation, some 450 meters in diameter flanked by cliffs, lying south of the rocky headland whose northernmost point is the site of the Point Arena lighthouse. The cove appears about one-third of the way above the lower edge of Figure 1 (based on the 2005 NAIP Imagery for USGS 7.5min quadrangle: 38123h6, available online at atlas.ca.gov/quads). Apart from a pier (better seen in Figure 2), raised high above water level on pilings and the focus of local commercial fishing and tourist activity, the waterfront area of Arena Cove is almost devoid of infrastructure though it lies within the city limits of Point Arena. This small city has a population of about 500, who mainly reside at elevations that place them above the level likely to be impacted by even the most severe tsunamis. Population is sparse both to the south and north of Point Arena but inundation of Manchester Beach State Park, north of the lighthouse, and the low-lying area near the mouth of the Garcia River (crossed by State Highway 1), needs consideration. A comprehensive study of potential tsunami inundation, for the entire California coastline, was conducted by the University of Southern California Tsunami Research Center. Funded through the California Emergency Management Agency (CalEMA), by the National Tsunami Hazard Program, the study (Barberopoulou et al, 2011) has produced a set of inundation maps for emergency planning purposes accessible online in various forms, including a tool MyHazard (myhazards.calema.ca.gov) enabling users to acquire information specific to their site of interest. The CalEMA inundation results are available in GIS form and those specific to the Arena Cove area are used throughout this report. In addition to underpinning the modeling effort, the digital elevation model (DEM) for the region, provided by the National Geophysical Data Center (NGDC), includes a 3-D oblique view that assists greatly in visualizing the study area. In Figure 3, the CalEMA inundation information is overlaid, together with descriptive labels on an extract from the NGDC image, available in full in the DEM Report (Friday et al., 2009).

Arena Cove, both in appearance (Figure 2) and population, has not changed substantially since the nineteenth and early twentieth centuries, when it was one of the “Dog Hole” ports of the Mendocino Coast (Haugan, 2005). So named for their small size, these ports nonetheless served an important role in the provision of lumber in the building of the cities of California, and in the rebuilding of San Francisco in the wake of the 1906 earthquake and fire (see the inset to Figure 2, reproduced by permission of the Mendocino County Historical Society.) The Point Arena lighthouse, and its namesake city, were seriously damaged by the earthquake and the San Andreas Fault (SAF) dominates the local topography. The SAF intersects the coast just north of Point Arena en route to the triple junction near Cape Mendocino, the southern limit of the Cascadia Subduction Zone, which constitutes a major earthquake and tsunami hazard to the U.S. west coast.

Unlike the other ports of the Mendocino Coast, whose mouths have sand bars, the channeling northward of the Garcia River by the SAF, leaves a limited watershed to

supply sediment to Arena Cove via Point Arena Creek entering the cove through a small, steep-sided valley. The current pier, rebuilt in 1986 following the damage to its predecessor by a series of storms in 1983, stands on pilings high above water level. It houses a crane to lower boats to the water and the instrumentation for the tide gage whose sensor is adjacent to the pier. In earlier days piers, and wire chutes atop the flanking cliffs, delivered lumber products to coastal schooners. Today's pier supports local commercial fishing and sightseeing, surfing and pier fishing are popular tourist activities.

Apart from the pier, and some riprap, the cove remains in its natural state. A congressional study (U.S. Secretary of War, 1914) considered the possibility of engineering works to make Arena Cove a "harbor of refuge" between San Francisco and Humboldt Bay but concluded that this was neither feasible nor a serious need. Consequently sea level data from the tide gage represents coastal conditions, unaffected by infrastructure. Port Road links the pier, the parking area, and some buildings housing fishing and tourist amenities, to the city proper. Accommodations, in the immediate pier area, are confined to an inn that is well elevated from the waves associated with winter storms and, as this report will document, even the most severe tsunami.

South of Arena Cove, as illustrated in a striking series of aerial photographs (www.californiacoastline.org the source of the main frame of Figure 2), high cliffs limit potential impact by tsunamis. To the north of the lighthouse however, and stretching as far as the Irish Beach community, lies Manchester Beach State Park. Inland from the point of entry to the ocean of the Garcia River, is the Manchester-Point Arena Rancheria of the federally recognized Band of Pomo Indians. While the historical record of tsunamis does not include mention of this area, its risk for inundation is evident in the CalEMA chart and the results of this study indicate that it may be prone to inundation in severe tsunami events. Thus, while the study focuses on Arena Cove, and the validation of the forecast model is provided by the tide gage there, the analysis of the most severe scenarios will consider potential impacts to the Manchester area. The community of Manchester itself appears to be immune to direct impact, though the State Highway 1 (also called the Shoreline Highway) may be inundated where it crosses the Garcia River. Queries to the CalEMA "MyHazards" site for Point Arena and Manchester show flooding and earthquake as other hazards to which they are prone, in addition to tsunami.

1.2. Natural Hazards

Several instances of mild tsunami signals are evident in the tide gage records for Arena Cove, whose name appears several times in the records compiled by Lander and Lockridge (1989) and the NGDC Tsunami Hazard Database (Dunbar, 2007; see www.ngdc.noaa.gov/hazard/). The historical record first mentions Mendocino County with a 1-meter wave height associated with the Sanriku event of 1896. O'Brien (1946) described a 2.4 meter wave (4.3m above MLLW) at Arena Cove during the 1946 Unimak tsunami. At Noyo Harbor (adjacent to Fort Bragg, the largest coastal community in the county; 2010 population 7,273) "100 fishing boats thrown 1.8m up beach and some damage to pier." While Arena Cove was not explicitly mentioned in connection with the 1957 Andreanof event there was a report from Noyo Harbor. Similarly during the 1960

Chile event, “6 boats broke mooring ... pier damaged” at Noyo Harbor and a height of 0.61m was observed at Gualala River near the southern boundary of Mendocino County. During the 1964 Alaska tsunami a run-up height of 1.83 meters occurred at Arena Cove and several instances of mild response to tele-tsunamis are available, following the installation of a tide gage in 1978, with which to validate model predictions.

The M_w 7.2 earthquake north of Cape Mendocino on April 25, 1992 was a very mild foretaste of a Cascadia Subduction Zone (CSZ) event. It produced wave heights of 0.14m at Arena Cove and 0.50m at Crescent City. Large-scale events on the CSZ are simulated later in the report but the weak 1992 event will be examined to see whether the presence of the Point Arena headland provides protection to Arena Cove, which lies in its lee for waves propagating along the coast from the north.

Combining events impacting northern California with those that have occurred since the Arena Cove tide gage was upgraded to 1-minute sampling, a total of 27 historical events are available for study. Nineteen of these, listed in Table 1a, are the standards for forecast model testing in the Pacific because their sea floor deformation is reasonably well known, either from the literature or, more recently, derived from direct observation of the wave trains they generated. The remaining eight, listed in Table 1b, have source characteristics that are less well known; they are included to expand the geographical coverage or because of their special relevance to Arena Cove. The Mendocino-1992 event for example was the most recent subduction-type event in Cascadia. Others, due to significant noise in the tide gage, do not produce a clear signal but shed light on Arena Cove as a reference point for coastal impacts. Figure 4 illustrates the distribution of the 27 historical sources. Those highlighted in red were employed for intercomparison of the reference and forecast versions of the model.

Direct seismic impact is another natural hazard to which Point Arena area is exposed. Its proximity to the rupture zone of the SAF in the San Francisco earthquake of 1906 resulted in significant damage to the town and the destruction of the lighthouse. While the SAF enters the ocean at Manchester Beach, its strike-slip nature reduces the likelihood of severe tsunami wave generation should ruptures occur in the immediate vicinity. Submarine landslides or collapse of sections of sea cliff are a potential local source for tsunami damage. Landslides triggered by seismic events caused significant loss of life during the 1929 Newfoundland event and accentuated the 1996 New Guinea tsunami. Landslide-generated tsunami waves are not currently included in the SIFT forecast methodology, nor are those generated meteorologically. However, to the extent that the waves they produce are detected by the DART[®] array, some warning of their presence may be available.

Table 1a. Source characterization for historical tsunami events employed in Arena Cove model testing. Those in bold text were used in RM/FM inter-comparison. Sources identified as “ad hoc” may not be identically defined in other Forecast Model reports.

Event	Earthquake / Seismic			Model		
	USGS Date Time (UTC) Epicenter	CMT Date Time (UTC) Centroid	Magnitude M _w	Tsunami Magnitude	Subduction Zone	Tsunami Source (Reference/Derivation)
1946 Unimak	01 Apr 12:28:56 52.75°N 163.50°W	Not Available	8.5	8.5	ACSZ	7.5×B23 + 19.7×B24 + 3.7×B25 (López & Okal, 2006)
1952 Kamchatka	04 Nov 16:58:26.0 52.76°N 160.06°E	Not Available	9.0	9.0	KISZ	19.71 × (A4 + Y4 + Z4 + A5 + Y5 + Z5 + A6 + Y6 + Z6) (ad hoc)
1957 Andreanof	09 Mar 14:22:31 51.56°N 175.39°W	Not Available	8.6	8.7	ACSZ	31.4×A15 + 10.6×A16 + 12.2×A17 (Preliminary)
1960 Chile	22 May 19:11:14 38.29°S 73.05°W	Not Available	9.5	9.5	CSSZ	125×(A93 + B93 + Z93 + A94 + B94 + Z94 + A95 + B95) Kanamori & Cipar (1974)
1964 Alaska	28 Mar 03:36:00 61.02°N 147.65°W	Not Available	9.2	8.9	ACSZ	15.4×A34 + 18.3×B34 + 48.3×Z34 + 19.4×A35 + 15.1×B35 (Tang et al., 2006, 2009)
1994 East Kuril	04 Oct 13:22:58 43.73°N 147.32°E	04 Oct 13:23:28.5 43.60°N 147.63°E	8.3	8.1	KISZ	9.0×A20 (ad hoc)
1996 Andreanof	10 Jun 04:03:35 51.56°N 175.39°W	10 Jun 04:04:03.4 51.10°N 177.410°W	7.9	7.8	ACSZ	2.40×A15 + 0.80×B16 (Preliminary)
2001 Peru	23 Jun 20:33:14 16.26°S 73.64°W	23 Jun 20:34:23.3 17.28°S 72.71°W	8.4	8.2	CSSZ	5.7×A15 + 2.9×B16 + 1.98×A16 (Preliminary)
2003 Hokkaido	25 Sep 19:50:06 41.77°N 143.904°E	25 Sep 19:50:38.2 42.21°N 143.84°E	8.3	8.3	KISZ	3.95 × (A22 + B22 + A23 + B23) (ad hoc)
2003 Rat Island	17 Nov 06:43:07 51.13°N 178.74°E	17 Nov 06:43:31.0 51.14°N 177.86°E	7.7	7.8	ACSZ	2.81×B11 (Real-time)
2006 Tonga	03 May 15:26:39 20.13°S 174.161°W	03 May 15:27:03.7 20.39°S 173.47°W	8.0	8.0	NTSZ	6.6×B29 (ad hoc)
2006 Kuril	15 Nov 11:14:16 46.607°N 153.230°E	15 Nov 11:15:08 46.71°N 154.33°E	8.3	8.1	KISZ	4.0×A12 + 0.5×B12 + 2.0×A13 + 1.5×B13 (Real-time)
2007 Kuril	13 Jan 04:23:20 46.272°N 154.455°E	13 Jan 04:23:48.1 46.17°N 154.80°E	8.1	7.9	KISZ	-3.64 × B13 (Real-time)
2007 Solomon	01 Apr 20:39:56 8.481°S 156.978°E	01 Apr 20:40:38.9 7.76°S 156.34°E	8.1	8.2	NVSZ	12.0×B10 (Preliminary)
2007 Peru	15 Aug 23:40:57 13.354°S 76.509°W	15 Aug 23:41:57.9 13.73°S 77.04°W	8.0	8.1	CSSZ	0.9×A61 + 1.25×B61 + 5.6×A62 + 6.97×B62 + 3.5×Z62 (Preliminary)
2007 Chile	14 Nov 15:40:50 22.204°S 69.869°W	14 Nov 15:41:11.2 22.64°S 70.62°W	7.7	7.6	CSSZ	1.65×Z73 (Real-time)
2009 Samoa	29 Sep 17:48:10 15.509°S 172.034°W	29 Sep 17:48:26.8 15.13°S 171.97°W	8.1	8.1	NTSZ	3.96×A34 + 3.96×B34 (Real-time)
2010 Chile	27 Feb 06:34:14 35.909°S 72.733°W	27 Feb 06:35:15.4 35.95°S 73.15°W	8.8	8.8	CSSZ	17.24×A88 + 8.82×A90 + 11.84×B88 + 18.39×B89 + 16.75×B90 + 20.78×Z88 + 7.06×Z90 (Real-time)
2011 Honshu	11 Mar 05:46:24 38.297°N 142.372°E	11 Mar 05:47:47.2 38.486°N 142.597°E	9.0	9.0	KISZ	4.66 × B24 + 12.23×B25 + 26.31×A26 + 21.27×B26 + 22.75×A27 + 4.98×B27 (Real-time; Tang et al., 2012)

Table 1b. Supplementary historical tsunami events employed for Arena Cove forecast model testing. Those identified as “ad hoc” and may not be identically implemented in other Forecast Model reports.

Event	Earthquake / Seismic			Tsunami Source (Reference/Derivation)		
	USGS Date Time (UTC) Epicenter	CMT Date Time (UTC) Centroid	M _w Magnitude	Tsunami Magnitude	Subduction Zone	Tsunami Source
1896 Sanriku	15 Jun 10:33:00 39.5°N 144.0°E		7.6	7.6	KISZ	b25 x 1.413 (ad hoc)
1992 Mendocino	25 Apr 18:06:04 40.368°N 124.316°W	25 Apr 18:06:11.8 38.56°N 123.31°W	7.2	7.2	ACSZ	a65 x 0.355 or b65 x 0.355 (ad hoc)
1995 Chile	30 Jul 05:11:24 23.340°S 70.294°W	30 Jul 05:11:56.9 24.17°S 70.74°W	8.0	8.0	CSSZ	2.812 x (a75 + b75) (ad hoc)
1995 Kuril	03 Dec 18:01:09 44.663°N 149.300°E	03 Dec 18:01:36.1 44.82°N 150.17°E	7.9	7.9	KISZ	1.991 x (a17 + z17) (ad hoc)
1996 Irian Jaya	17 Feb 05:59:31 0.891°S 136.952°E	17 Feb 06:00:02.8 0.67°S 136.62°E	8.2	8.2	NGSZ	2.7984 x (a9 + b9 + a10 + b10) (ad hoc)
2009 PapuaNG	03 Jan 19:43:51 0.414°S 132.885°E	03 Jan 19:44:09.0 0.38°S 132.83°E	7.6	7.6	NGSZ	0.7046 x (b13 + b14) (ad hoc)
2009 Kuril	15 Jan 17:49:39 46.857°N 155.154°E	15 Jan 17:49:48.3 46.97°N 155.39°E	7.4	7.4	KISZ	b12 x 0.7063 (ad hoc)
2009 Vanuatu / Santa Cruz	07 Oct 22:03:15 13.052°S 166.187°E	07 Oct 22:03:28.9 12.59°S 166.27°E	7.6	7.6	NVSZ	1.2xB24 + 0.26xA23 followed after 15minutes by 2.6xB23 + 0.9xA23
	07 Oct 22:18:26 12.554°S 166.320°E	07 Oct 22:19:15.3 11.86°S 166.01°E	7.8	7.9	NVSZ	(Preliminary Wei 2009, Personal Communication)

Another local hazard that has been a frequent cause of damage to Arena Cove has been ocean wave action. Originating locally, or as swell from distant storms, such waves caused severe damage to the pier in 1983 that necessitated its replacement. Another impact of ocean waves, of relevance to tsunami detection and modeling, is in the noise they produce in the tide gage record that can mask weaker tsunami signals, Harbor resonance in the case of Crescent City can amplify the tsunami and may be a factor too in the Arena Cove response.

1.3. Tsunami Warning and Risk Assessment

The forecast model development, described here, will permit Arena Cove, CA, to be incorporated into the tsunami forecasting system SIFT, developed at NCTR (NOAA Center for Tsunami Research) and now in operational use at the U.S. Tsunami Warning Centers (TWC's). The system has had considerable success in accurately forecasting the impact of both moderate and severe tsunami events in recent years and in the following section the methodology that permits such forecasts is discussed as prelude to a description of development of the forecast model for Arena Cove. With the model in hand, validated with historical events and with its stability verified by extensive testing against extreme scenarios, real-time forecasts will be available to inform local emergency response. Additionally, the synthetic scenarios investigated during model development, and reported here, provide an initial tsunami risk assessment as described in the Results and Discussion section.

2.0 Forecast Methodology

2.1 The Tsunami Model

In operational use, a tsunami forecast model is used to extend a pre-computed deep-water solution into the shallows, and onshore as inundation if appropriate. The model consists of a set of three nested grids, of increasingly fine resolution that, in a real-time application of the MOST model (Method of Splitting Tsunami: Titov and Synolakis, 1998; Titov and González, 1997), permits forecasts at spatial scales (as little a few tens of meters) relevant to local emergency management. The validity of the MOST model applied in this manner, and the operational effectiveness of the forecast system built around it, has been demonstrated during unplanned tests triggered by several mild to moderate tsunami events in the years since the 2004 Indian Ocean disaster (Wei et al., 2008). Successful hindcasting of observed historic events, even mild ones, during forecast model development lends credence to the ability to accurately forecasting the impact of future events. Such validation of tsunami modeling procedures is documented in other volumes of the series of which this report is but one. Before proceeding to a description of the forecast model development for Arena Cove, it is useful to describe the steps in the overall forecast process.

2.2 The SIFT Forecast System

Operational tsunami forecasts are generated at Tsunami Warning Centers, staffed 24/7 in Alaska and Hawaii, using the SIFT (Short-term Inundation Forecasting for Tsunamis) tool, developed at NCTR. The semi-automated process facilitates the steps by which TWC operators assimilate data from an appropriate subset of the DART® tsunami sensors, “invert” the data to determine the linear combination of pre-computed propagation solutions that best match the observations, then initiate a set of forecast model runs if coastal communities are threatened or, if warranted, cancel the warning.

Steps in the process are as follows:

- When a submarine earthquake occurs the global network of seismometers registers it. Based on the epicenter, the unit sources in the Propagation Database (Gica et al., 2008) that are most likely to be involved in the event, and the DART® array elements (Spillane et al., 2008) best placed to detect the waves passage are identified. TWC watch-standers can trigger DART®s into rapid sampling mode in the event that this did not occur automatically in response to the seismic signal.
- There is now an unavoidable delay while the tsunami waves are in transit to the DART®s; at least a quarter of a cycle of the first wave in the train must be sampled before moving to the “inversion” step.
- When sufficient data have accumulated, at one or more DART®s, the observed time series are compared with the model series from the candidate unit sources. Since the latter are pre-computed (using the MOST code), and the dynamics of tsunami waves in deep water is linear, a least squares approach taking very little time can identify the unit sources, (and the appropriate scale factors for each,) that

best fit the observations. The “inversion” methodology is described by Percival et. al., (2009).

- Drawing again on the Propagation Database, the scale factors are applied to produce a composite basin-wide solution with which to identify the coastal regions most threatened by the radiating waves.
- It is at this point that one or more forecast models are run. The composite propagation solution is employed as the boundary condition to the outermost (A-grid) domain of a nested set of three real-time MOST models that telescope with increasingly fine scale to the community of concern. A-grid results provide boundary conditions to the B-grid, which in turn forces the innermost C-grid. Non-linear processes including inundation are modeled so that, relying on the validation procedures during model development, credible forecasts of the current event are available.
- Each forecast model provides quantitative and graphic forecast products with which to inform the emergency response, or to serve as the basis for canceling or reducing the warnings. Unless the tsunami source is local, the forecast is generally available before the waves arrive but, even when lead-time cannot be provided, the several hour duration of a significant event (in which the first wave may not be the most damaging) give added value to the multi-hour forecasts provided.

Because multiple communities may be potentially at risk, it may be necessary to run simultaneously, or in a prioritized manner, multiple forecast models. Each must be optimized to run efficiently in as little time as possible; the current standard is that an operational forecast model should be capable of simulating 4 hours of real time within about 10 minutes of CPU time on a fast workstation computer.

3.0 Model Development

3.1 Digital Elevation Models

Water depth determines local tsunami wave speed and sub-aerial topography determines the extent to which tsunami waves inundate the land. Thus a prerequisite for credible tsunami modeling is the availability of accurate gridded bathymetric and topographic datasets, termed DEM's (Digital Elevation Models.) Given their expertise in this area, and the number of coastal communities needing tsunami forecast capability, NCTR relies heavily on the National Geophysical Data Center (NGDC) to provide the DEM's needed. In the case of Arena Cove, the DEM, a composite of multiple data sources merged and converted to a common datum of Mean High Water (MHW), was produced and documented by Friday et al. (2009). The use of MHW as the "zero level" for forecast results is standard. The MOST model does not include tidal fluctuations and, since a tsunami may arrive at any stage of the tide, it is best to employ a "worst-case" approach by assuming high tide when forecasting inundation. For some Forecast Models grounding of vessels and the strong and the rapidly varying currents often associated with even mild tsunamis are of concern. For Arena Cove, lacking a marina and shoreline infrastructure, low water impacts are less important.

Table 2. The main features of the Arena Cove Digital Elevation Model (DEM).

Grid Area	Arena Cove, California
Coverage Area	123.43° to 124.43° W; 38.40° to 39.40° N
Coordinate System	Geographic decimal degrees
Horizontal Datum	World Geodetic System 1984 (WGS84)
Vertical Datum	Mean High Water (MHW)
Vertical Units	Meters
Cell Size	1/3 arc-second
Grid Format	ESRI Arc ASCII grid

The DEM provided by NGDC for the Arena Cove area was illustrated in Figure 3; its salient features listed in Table 2 reproduced from DEM documentation (Friday et al., 2009). The NGDC report thoroughly describes the data sources and methods employed in constructing the DEM. With one-third arc second (~10m) resolution, the DEM provides the basis for the B and C-grids for both reference and forecast model usage. NCTR maintains an atlas of lower resolution gridded bathymetries, which can be used for the A-grids, as described later. All of the DEMs employed were verified for consistency with charts, satellite imagery, and other datasets during the course of MOST grid development.

The bathymetry and topography used in the development of this forecast model was based on the digital elevation model provided by the National Geophysical Data Center and the author considers it to be a good representation of the local topography/bathymetry. As new digital elevation models become available, forecast models will be updated and report updates will be posted at http://nctr.pmel.noaa.gov/forecast_reports/.

3.2 Tides and Sea Level Variation

Arena Cove's history of tidal observations dates back only to 1978. The tide station (9416841, 38°54.8'N, 123°42.4'W) is located near the end of the pier whose concrete pilings raise the deck about 25 feet above sea level and do not impede water movement within the cove. The instrumentation was upgraded in 2006 to include a tsunami-capable gage sampling at 1-minute intervals; some earlier data was sampled at 6-minute intervals and several historical events are only available as marigrams on microfiche. An ongoing project at NGDC will digitize the more critical images in this archive.

Station characteristics for 9416841 are provided in Table 3, based on the wealth of online tidal information available at NOAA's CO-OPS (Center for Operational Oceanographic Products and Services) website (tidesandcurrents.noaa.gov). Note the sizeable diurnal range of over three meters and that, while the long-term rate of change in sea level is low (compared to more seismically active areas), there is substantial seasonal, interannual and short-term variability. Owing to the relative short history of the Arena Cove gage, trends and cycles are reported for Crescent City to the north and Point Reyes to the south.

Table 3. Tidal characteristics of the Arena Cove, CA Tide Gage (9416841).

Arena Cove, CA		Station#9416841	38°54.8'N, 123°42.4'W	
Tidal Datum and Range Values (Epoch 1983-2001)				
MHHW (Mean Higher High)	10.618m	Great Diurnal Range 1.792m	Mean Range 1.233m	
MHW (Mean High Water)	10.413m			
MSL (Mean Sea Level)	9.786m			
MLW (Mean Low Water)	9.180m			
MLLW (Mean Lower Low)	8.826m			
Sea Level Trends and Cycles (from Point Reyes, CA #9415020)				
Long Term SL Trend	Increasing 2.10±1.52mm/year			
Seasonal Cycle Range	Minimum -89mm(April); Maximum 59mm(September)			
Interannual Variation (from1980)	Minimum -20mm(1988); Maximum +21mm(1997)			
Sea Level Trends and Cycles (from Crescent City, CA #9419750)				
Long Term SL Trend	Decreasing 0.65±0.36mm/year			
Seasonal Cycle Range	Minimum -87mm(May); Maximum 85mm(January)			
Interannual Variation (from1980)	Minimum -20mm(1989); Maximum +28mm(1998)			

A sample section of the tide gage record, again extracted from the CO-OPS website is reproduced in Figure 5. Deviations (or residuals) from the astronomically predicted tide can be several centimeters and the variability strong. In particular the highest water level reported for the Arena Cove gage is 1.048m above MHW (Feb 6, 1998) so the use of MHW as the zero level of modeled sea level may underestimate the truly worse case.

While the simultaneous arrival of the crest of a large tsunami at high tide during a storm surge has low probability, a feature of the simulated events reported below is that sustained oscillations at a resonant period may extend the duration of the threat. This effect is notorious at Crescent City, CA which is frequently the most heavily impacted U.S. west coast location for remote events.

3.3 The CFL Condition and other considerations for grid design

Water depth dependent wave speed, in conjunction with the spacing of the spatial grid representation, place an upper limit on the time step permissible for stable numerical solutions employing an explicit scheme. This is the CFL limit (Courant-Friedrichs-Levy), which requires careful consideration when the grids employed for a reference or forecast model are being designed. Finer-scale spatial grids, or greater water depths, require shorter time steps thereby increasing the amount of computation required to simulate a specific real time interval.

Another feature of the application of gridded numerical solutions to the tsunami wave problem is the shortening that the wave train encounters in moving from deep water onto the shelf. In deep water a grid spacing of 4 arc-seconds (of latitude and longitude, corresponding to ~7km) is normally used to represent propagating wave trains whose wavelength is typically of the order of a few hundred kilometers. The stored results of such propagation model runs are typically decimated by a factor of 4, resulting in a database of ~30km spacing (and 1 minute temporal sampling) with which to generate the boundary conditions for the outermost of the nested grids in a model solution. The extraction of the boundary conditions (of wave height and the two horizontal velocity components) is achieved by linear interpolation in space and time. To provide realistic interpolated values the stored fields for these variables must be smoothly varying, and have adequate sampling in space and time to resolve their structure. This necessitates the placement of the offshore boundary of the forecast model domain well offshore. The presence of the Mendocino Escarpment is another incentive to do so, in order that its role in topographic steering of trans-Pacific wave trains be adequately represented.

Figure 6 illustrates the placement of the model domain in its west coast setting. The outermost A-grid covers the entire region shown; embedded in it is the B-grid, which covers most of Mendocino County. The innermost C-grid, with the finest spatial resolution spans the region north and south of Point Arena. Marked in red are a number of nearby communities where run-ups are mentioned in the historical record. Indicated as black triangles are the tsunami-capable tide gages of the region, the closest of which are Point Reyes to the south and North Spit to the north. Almost directly offshore is DART® 46411. This would play a major role in the detection of regionally generated waves. Its offshore location cleanly registers moderate to large tele-tsunamis and could, potentially, refine a local forecast that was initially based on DART® array elements closer to the source. Red, green, and magenta lines indicate (using the color-convention employed in the online USGS/NEIC earthquake resources) the three types of fault that radiate from the triple junction off Cape Mendocino. To the south is the strike-slip San Andreas Fault, skirting the coastline north of San Francisco Bay before entering the ocean within the C-

grid domain. The Mendocino Escarpment is dramatic evidence of the ridge fault extending offshore but of most concern as a local source of tsunamis is the Cascadia Subduction Zone. The two southernmost pairs of the unit source set used to represent it fall within the A-grid domain. A “beachball,” that visually represents the source mechanism, marks the location of the Mendocino-1992 event which was the last significant subduction event in Cascadia.

3.4 Specifics of the model grids

After several rounds of experimentation, the extents and resolutions of the nested grids were chosen, and are illustrated in Figures 7 and 8; details are provided in Tables 4 and 5. The Reference Model (RM) and Forecast Model (FM) grid pairs (at the A and B levels) have the same extent, differing only in resolution; the C-grid domain is however slightly larger for the RM than for the FM, the dimensions of the latter being reduced to achieve a shorter run-time appropriate to operational use. The corresponding panels in the figures employ the same depth contours and color palette, which consequently are only shown in the FM version (Figure 8). Rectangles drawn in red for the A and B grid panels, indicate the extent of the embedded grid; where appropriate the blue rectangles indicate the less extensive FM C-grid. Superimposed in the C-grid panels is the network of rivers, creeks and roads. The thick red line marks State Hwy 1, also called the Shoreline Highway.

Table 4. Specifics of the grids and model parameters employed to model Arena Cove, CA. “EWxNS” denotes the number of grid values in the zonal (East to West) and meridional (North to South) directions respectively.

Grid	Zonal Extent (W)		Meridional Extent (N)		Resolution (“)		Values EW x NS	
					RM	FM	RM	FM
A	128.00 ⁰	121.50 ⁰	36.00 ⁰	42.50 ⁰	30	60	781x781	391x391
B	124.55 ⁰	123.00 ⁰	38.35 ⁰	39.80 ⁰	6	24	931x871	234x291
C	123.85 ⁰	123.60 ⁰	38.82 ⁰	39.03 ⁰	1	2	901x757	235x313
	123.78 ⁰	123.65 ⁰	38.89 ⁰	39.02 ⁰				

Table 5. Grid file names and grid-related parameters. The time steps for the A and B-grids must be integer multiples of the basic time step chosen for the C-grid.

Grid	Filename	Maximum Depth (m)	Minimum CFL (s)	Model Time Step (s)	Water Cells
A	ArenaCoveCA_RM_A	5002	3.350	3.0 (5x)	436966
	ArenaCoveCA_FM_A	5005	6.689	6.0 (4x)	109323
B	ArenaCoveCA_RM_B	3781	0.7559	0.6 (1x)	485760
	ArenaCoveCA_FM_B	3776	2.893	1.5 (1x)	40535
C	ArenaCoveCA_RM_C	143.6	0.6423	0.6	434701
	ArenaCoveCA_FM_C	94.4	1.526	1.5	37611

Both C-grids lie entirely within the NGDC-provided Arena Cove DEM; A and B-grids incorporate bathymetric and topographic from other DEM datasets available at NCTR. Some smoothing and editing were necessary to eliminate erroneous points or grid features that tend to cause model instability. For example, “point” islands where an isolated grid cell stands above water are eliminated, as are narrow channels or inlets one grid unit wide; these tend to resonate in the numerical solution. Large depth changes between adjacent grid cells can also cause numerical problems; customized tools (such as “bathcorr”) are available to correct many of these grid defects.

Details of the model grids are provided in Tables 4 and 5. The latter lists the maximum depth, the CFL time step requirement that must not be exceeded, and the actual time steps chosen for the reference and forecast model runs. Since in the current version of MOST, employed by SIFT, the numerical solutions in the three grids proceed simultaneously, there is a requirement that the A and B-grid time steps be integer multiples of the (innermost) C-grid time step in addition to satisfying the appropriate CFL requirement. For both reference and forecast models the CFL requirement of the C-grid was the most stringent. The values chosen are shown in the final column of Tables 4 and 5 and are such that an integer multiple of each time step (16x for the forecast model, 64x for the reference) is identically 30 seconds, the chosen output time interval for both models. When run on an Intel® Xeon® E5670 2.93GHz processor, the forecast model can simulate four hours in 10.73 minutes, close to the desired value of 10-minutes for this metric.

3.5 Model Run Input and Output Files

In addition to providing the bathymetry file names and the appropriate time step and A, B grid multiples as provided in the tables above, the designer must provide a number of additional parameters in an input file. These include the Manning Friction Coefficient, a depth threshold to determine when a grid point becomes inundated, and the threshold amplitude at the A-grid boundary that will start the model. An upper limit is specified in order to terminate the run if the wave amplitude grows beyond reasonable expectation. Standard values are used: 0.0009 for the friction coefficient and 0.1m for the inundation threshold. The latter causes the inundation calculation to be avoided for insignificant water encroachments that are probably below the uncertainty in the topographic data. Inundation can, optionally, be ignored in the A and B-grids, as is the norm in the (non-nested) MOST model runs that generate the propagation database. When A and/or B-grid inundation is excluded, water depths less than a specified “minimum offshore depth” are treated as land; in effect a “wall” is placed at the corresponding isobath. When invoked, a value of 5m is applied as the threshold, though A and B inundation is normally permitted as a way to gain some knowledge of tsunami impact beyond the scope of the C-grid domain. Other parameter settings allow decimation of the output in space and/or time. As noted earlier, 30-second output has been the target and output at every spatial node is preferred. These choices avoid aliasing in the output fields that may be suggestive of instability (particularly in graphical output), when none in fact exists.

Finally the input files (supplied in Appendix A for both the reference and forecast versions of the model) provide options that control the output produced. Output of the

three variables: wave amplitude, and the zonal (positive to the east) and meridional (positive to the north) velocity components can be written (in netCDF format) for any combination of the A, B, and C-grids. These files can be very large! A separate file, referred to as a “SIFT” file, contains the time series of wave amplitude at each time step at discrete cells of a selected grid. Normally the time series at a “reference” or “warning point”, typically the location of a tide gage is selected to permit validation in the case of future or historical events. Also output in the SIFT file is the distribution of the overall minimum and maximum wave amplitude and speed in each grid. By contrast with the complete space-time results of a run, the SIFT file (also netCDF) is very compact and, if more than a single grid point is specified, a broader view of the response is provided.

By default two additional output files are generated: a listing file, which summarizes run specifications, progress, and performance in terms of run time. Also included in this file is information to determine the reason, should a run not start or terminate early. Finally a “restart” file is produced so that a run can be resumed, beginning at the time it ended, either normally or by operator intervention.

The input files described above are specific to the model itself. For an actual run, the program must be pointed toward the files that contain the boundary conditions of wave amplitude (HA), and velocity components (UA, VA), to be imposed at the A-grid boundary. Time varying conditions are generally extracted as a subset of a basin-wide propagation solution (either a single unit source or several, individually scaled and linearly combined) that mimic a particular event. These boundary-forcing files typically consist of 24 hours of values (beginning at the time of the earthquake), sampled at 1-minute intervals and available on a 16 arc-minute grid. Occasionally, for more remote seismic sources (or when delayed arrival of secondary waves due to reflections are a concern, as has been seen at Hawaii,) the time span of the propagation run available for forcing is extended beyond one day.

4.0 Results and Discussion

Before proceeding to an extensive suite of model runs, that explore the threat to the Point Arena area from various source regions, the stability of the model is tested in both low and extreme amplitude situations. The former we refer to as “null source” testing: where the boundary forcing is at such a low level (but not precisely zero of course) that the response is expected to be negligible. These tests can be highly valuable in revealing localized instabilities that may result from undesirable features in the discretized bathymetric representation. Inlets or channels that are only one grid cell wide may “ring” or resonate in a non-physical way in the numerical solution. An instability may not grow large enough to cause the model to fail but, in a run with typical tsunami amplitudes, may be masked by actual wave variability.

Forcing by extreme events should also be tested. In addition to the need to test model stability under such circumstances, there is a parameter in the input file that truncates the run if a prescribed threshold is exceeded. For operational use, the threshold must be set high enough so that an extreme event run is not unnecessarily terminated. Both tests should be done for test sources whose waves enter the model domain from different directions since, although stable for one set of incoming waves, an instability may be encountered for another. The “null” and “extreme” testing of the forecast and reference models is reported in the following subsections. Further evidence of stability is provided by the extensive set of scenarios, aimed at exploring the dependence of impact to source location, described later in the report, and in independent testing by other members of the NCTR team before the model is released for operational use.

4.1 The “Null” Tests

Three null test cases (see Table 6) were run representing sources in the western Aleutians, the Philippines, and south of Japan. Based on sources from the propagation database (Gica et al., 2008), their amplitudes were scaled down by a factor of 10,000 so as to mimic an $M_w=4.8333$ / Slip 0.0001m source rather than the $M_w=7.5$ / Slip 1m standard. A number of grid cells in the B and C grids emerged as potential sources of instability. These were generally minor indentations of the coastline, barely resolved by the grids, or narrow channels. Also to be looked for in further testing is the area northwest of the Point Arena Light when the rugged seabed reveals several past water level stands. A limited number of grid cells in the outermost (A) grid required correction. Generally these were associated with non-physical features in the topographic database, such as where a track of ship-based soundings were improperly merged with other data sources. After an iterative process of grid correction and retesting using these “null” sources, both of the reference (RM) and forecast model (FM) grids were deemed satisfactory and the testing of realistic events can begin. Figure 9 illustrates a step in the process where a deficiency in the RM grid generated a mild instability (in the EPSZ B19 micro-tsunami scenario – see Table 6) causing the RM time series at the reference point, initially in close agreement with the FM, to develop unrealistic, high frequency oscillations. Though still generally tracking the FM result, and not growing without bound, the feature could behave erratically in simulating real events. Modification of the

Table 6. Synthetic tsunami events employed in Arena Cove, CA model testing. The RM and FM solutions of those shown in bold text were inter-compared extensively.

Scenario Name	Source Zone	Tsunami Source	α [m]
Mega-tsunami (M_w 9.3) Scenario			
KISZ 1-10	Kamchatka-Yap-Mariana-Izu-	A1-A10, B1-B10	25
KISZ 22-31	Kamchatka-Yap-Mariana-Izu-Bonin	A22-A31, B22-B31	25
KISZ 32-41	Kamchatka-Yap-Mariana-Izu-Bonin	A32-A41, B32-B41	25
KISZ 56-65	Kamchatka-Yap-Mariana-Izu-Bonin	A56-A65, B56-B65	25
ACSZ 6-15	Aleutian-Alaska-Cascadia	A6-A15, B6-B15	25
ACSZ 16-25	Aleutian-Alaska-Cascadia	A16-A25, B16-B25	25
ACSZ 22-31	Aleutian-Alaska-Cascadia	A22-A31, B22-B31	25
ACSZ 50-59	Aleutian-Alaska-Cascadia	A50-A59, B50-B59	25
ACSZ 56-65	Aleutian-Alaska-Cascadia	A56-A65, B56-B65	25
CSSZ 1-10	Central and South America	A1-A10, B1-B10	25
CSSZ 37-46	Central and South America	A37-A46, B37-B46	25
CSSZ 89-98	Central and South America	A89-B98, B89-B98	25
CSSZ 102-111	Central and South America	A102-A111, B102-	25
NTSZ 30-39	New Zealand-Kermadec-Tonga	A30-A39, B30-B39	25
NVSZ 28-37	New Britain-Solomons-Vanuatu	A28-A37, B28-B37	25
MOSZ 1-10	ManusOCB	A1-A10, B1-B10	25
NGSZ 3-12	North New Guinea	A3-A12, B3-B12	25
EPSZ 6-15	East Philippines	A6-A15, B6-B15	25
RNSZ 12-21	Ryukus-Kyushu-Nankai	A12-A21, B12-B21	25
M_w 7.5 Scenario			
NTSZ B36	New Zealand-Kermadec-Tonga	B36	1
Micro-tsunami Scenario			
EPSZ B19	East Philippines	B19	0.01
RNSZ B14	Ryukus-Kyushu-Nankai	B14	0.01
ACSZ B6	Aleutian-Alaska-Cascadia	B6	0.01

RM bathymetry eliminated the problem, as seen in the lower panel, and “null” tests involving other sources (RNSZ B14 and ACSZ B6) did not reveal other issues.

4.2 The Extreme Case Tests

The record of tsunami impact on the northern California coast discussed later reveals that sources around the entire periphery of the Pacific can be felt. Indeed the catastrophic Indian Ocean tsunami of 2004 was detectable at Arena Cove as it was throughout the global ocean. A broad suite of 19 extreme events (so-called mega-tsunamis) whose locations are standard for testing of Pacific basin forecast models, are described in Table 6. Their locations are shown in Figure 10. To simulate each mega-tsunami source, ten A-B pairs of unit sources are used, with an evenly distributed slip of 25m. As described by Gica et al. (2008), each unit source represents a 100x50km area of the fault surface with the long axis parallel to the plate boundary. The B-row is shallowest, sloping from a nominal depth of 5km (unless a depth estimate has been provided by the USGS based on the earthquake catalogs), row-A is deeper, followed by rows Z, Y, X, ... where appropriate. Thus, the extreme case sources represent 1,000 km long ruptures with a width of 100km; the corresponding magnitude is $M_w=9.3$.

Discussion of the entire set in greater detail is provided later in the report, once the validity of the Forecast Model has been established. Here we focus on a subset of three, highlighted in Figure 10 and Table 6, to contrast the Forecast Model (FM) with the more highly resolved Reference Model (RM). The results are presented in Figures 11-13, with the time series at the reference point (the Arena Cove tide gage) shown in the upper panel and the amplitude and current pattern at a selected time shown below. The black curve and red curves represent the RM and FM respectively; the green line identifies the time at which the comparison in the lower panel was made. Inset in the lower panels are enlargements of the area around Arena Cove and are left pixilated to reflect the discrete grid resolution.

It is noticeable that, in all three of the cases shown, the RM tends to oscillate longer and have somewhat larger amplitude than does the FM though the two solutions are in close agreement for the first few tsunami waves. This is likely a physical reality: the more highly resolved bathymetry and coastline of the RM providing greater scope for non-linear features or reflected waves to develop. This observation suggests a caveat to operational use of the FM: while accurate portrayal of the early history of an event is to be expected, the duration of the event and the amplitude of later waves may be underestimated. Tide gage data will be needed to verify this conjecture, which is pursued later in the report.

The snapshot comparisons in the lower panels of Figures 11-12 are quite reasonable, illustrating that the solutions match not just at the reference point. It is worth noting too that, although the ACSZ 56-65 mega-event represents a massive Cascadia tsunami, the scale of the impact to the Arena Cove area (~ 3m) is not substantially greater than from trans-Pacific locations (KISZ 01-10 off Kamchatka and NTSZ 30-39 near Samoa.) The Crescent City response to the same synthetic Cascadia mega-event exceeds 10m (Arcas

and Uslu, 2010). It would appear that the energy propagated along shore to the south, perhaps with some sheltering by Cape Mendocino, is reduced and that perhaps the greatest impact to Arena Cove may be associated with source regions elsewhere in the Pacific basin.

In Figure 13 the comparison time was intentionally chosen later in the event as a counterexample. While the reference point amplitudes and nearby fields the FM and RM may be in reasonable agreement, the broader wave patterns may have substantial phase differences. The comparisons in these lower panels is restricted to the portion of C-grid area common to both models. There is a suggestion that the near shore velocity fields at the north and south FM boundaries differ somewhat from the RM for which these are internal points.

Before proceeding to validate the model with historical events, one other synthetic event is usual in the testing protocol: a mild source of magnitude 7.5 at a remote location. A single unit source near Samoa (NTSZ-B36) is employed and its representation by the RM and FM are compared in Figure 14. Such an event results in a response of about 2cm in Arena Cove sea level and there is excellent agreement between both model representations in the earlier portion of the event.

Overall, the close agreement between the first wave arrival time and waveform, and overall range of variation of the two model representations in synthetic scenarios (even though the amplitude and phase is not always well-matched for later waves) suggests that the forecast model is performing well, and that we can, with confidence, proceed to model real events.

4.3 Model Validation with Historical Events

We now proceed to examine how well the RM and FM solutions compare with observation for several historical cases: those highlighted in Table 1 and Figure 6. Since the observations are limited to the tide gage records or run-up reports in Arena Cove the purpose of the lower panels is only to illustrate the agreement between the models.

The results, displayed and described below, represent the large Unimak-1946 and Alaska-1964 events, and three more recent ones: Kuril-2006 (which has been extensively studied), Samoa-2009, and Chile-2010. The latter three occurred subsequent to the installation of an improved tide gage at Arena Cove. In a later subsection, the Honshu tsunami of March 11, 2011 is discussed. It occurred while this report was undergoing internal review at NCTR but the FM was available for use in real-time circumstances. There is another difference between the earlier and more recent events. Source characterization for the former is based on the literature with the source mechanism estimated from the seismic record. The Kuril-2006 event was the first substantial event for which direct observation of the tsunami wave train was available from multiple deep-water DART® sites. As such its source characteristics, and those for Samoa-2009, Chile-2010, and Honshu-2011 are better suited to tsunami modeling and forecast; those based on seismic data only may suffer from the defect that earthquakes differ in their ability to

generate tsunami waves. An extreme case of this is the Sanriku-1896 event, which is modeled and discussed briefly later in this report. It was a so-called “tsunami-earthquake” (Dudley and Lee, 1998), causing devastating losses in Japan despite its modest magnitude and scant warning in the form of ground motion.

Even in the case of source characterizations based on DART® detection and inversion, it should be borne in mind that perfect agreement between the model wave and observation is unlikely. For one thing, the DART® sites used in the inversion process may be well described by a linear combination of unit source functions, but their placement may limit the ability to predict basin-wide energy propagation. Ideally one might hope to refine the model solution in light of DART® observations closer to the impact site. The deep-water waves in the far field (for example 46411 in the case of Arena Cove) may however fall below the DART® detection threshold. Neither are the tide-gage observations, available for comparison with model prediction, perfect. They may include noise, possible amplified by harbor resonances and wind wave activity.

The Unimak-1946 and Alaska-1964 events were widely felt along the U.S. west coast, though the greatest impact was to the Hawaiian Islands. Reported run-ups at Arena Cove were 2.40 and 1.83 meters respectively, comparable in the case of Alaska-1964 (but somewhat lower for Unimak-1946), to the modeled responses shown in Figures 15 and 16. The RM and FM solutions match well, both in the time series in the upper panel and the amplitude and velocity field at the selected comparison time.

For the more recent events, where time series at Arena Cove permit direct inter-comparison with the RM and FM predictions, the results are presented in Figures 17-19. For the Kuril-2006 event, the reported 61cm run-up at Arena Cove exceeds, by a factor of about 2, the amplitude of the tide gage oscillations. Particularly for the early waves the model gives a reasonable representation of both the amplitude and timing of the observations. The time axis is in model hours and the discrepancy in the first wave arrival time is about 5 minutes, just 1% of its transoceanic travel time.

For Samoa-2009 the reported run-up at Arena Cove is 44cm, which may correspond to later in the record when harbor resonances may have been excited. For the early waves, the amplitude of the observations is closer to 20cm and, though it does reasonably well in predicting the early timing and the sequence of waves, the model underestimates the amplitude by about a third. For the Chile-2010 event the amplitude of the observations is replicated more closely. Again though, the reported run-up of 35cm is substantially greater than the greatest positive excursion of the de-tided observations as displayed.

Considering the above results, the main discrepancy appears to be the mismatch between reported run-up and the processed sea level time series. Some possible explanations come to mind. Run-up is defined as maximum elevation above the predicted tide, which may not include seasonal or meteorologically driven departures, which, as illustrated in Figure 5, can be several centimeters. Another possibility is that the overall maximum of the tide gage record may be aliased by high-frequency variability, which was smoothed somewhat by a 3-point running average in the preparation of these graphics.

4.4 Further Historical Simulations

The above analysis has documented good agreement between the forecast model and the slower running reference version. This permits us to simulate the balance of the historical cases where impacts to Arena Cove and northern California have been reported, and the remaining mega-tsunami scenarios with the forecast model alone. These runs are intended to further validate the stability of the FM but also provide some information on the exposure of the region to tsunamis generated at various points on the periphery of the Pacific.

In Figures 20-24 the full set of observed records at Arena Cove (or in some cases proxy sites) are compared with FM prediction. Also provided, for each event, is the state of the tide at Arena Cove. While probably of little concern for weak events, this may be a factor in the impact of larger ones. Reported run-up is included in each case though, as noted earlier, this may be only loosely related to the plotted series. In each case, the FM series is shown in black; the observations are drawn in red. Although studies of the global ocean response to the Indian Ocean tsunami of December 2004 suggest a run-up of 19 cm in Arena Cove, the signal is largely obscured by noise. An attempt was made to employ global ocean model results (on a coarser grid than is available for the Pacific propagation database) to drive the Arena Cove FM; the results were unsatisfactory and will not be presented. When a better resolved global solution is available, this event may be added to the suite employed for forecast model testing since it should shed light on the extent to which bathymetric resolution may impact arrival time accuracy.

The sequence begins with a cautionary tale: the “tsunami-earthquake” induced Sanriku-1896 event. This was modeled by a suitably positioned unit source (KISZ-B25) with the slip appropriate to the reported $M_w=7.6$ magnitude. As shown, such an event would be expected to generate only a few centimeter signal at Arena Cove. Large run-ups, one meter in the case of the nearby town of Mendocino, occurred illustrating the fact that direct observation of deep-water waves is needed for realistic forecasting. The depth and frequency of sea floor motion for this event was such that the earthquake magnitude poorly indicated its devastating tsunami-generating potential to Japan’s Sanriku coast.

Next consider the set of events from 1946-1964 that were felt in or near Arena Cove, though a tide gage had not yet been installed and the DART® array was still in the future (Unimak-1946 and Alaska-1964 were shown earlier.) In each case the source was represented by a weighted group of unit sources from the propagation database, or constructed to match source characteristics appearing in the literature (see Table 1 and Tang et al., 2006).

A number of other events between 1994 (East Kuril) and 2003 (Rat Island), listed in Table 1 and illustrated in Figures 21 and 22, generated weak responses in Arena Cove. In the case of East Kuril-1994, although the match is quite good, the presence of substantial noise in the tide gage record in advance of the waves’ arrival, suggests a limitation on the detection of weak tsunami signals. Particularly in the winter months the tide gage record

at Arena Cove can be extremely noisy. This is true for the IrianJaya-1996 event; for Chile-1995 and Kuril-1995 the tide gage records are not readily available though they may be digitized in the course of an NGDC project. For the Andreanof-1996 event the model seems to capture the timing and periodicity of the Arena Cove response as it does perhaps for the Rat Island-2003 event. For the Peru-2001 and Hokkaido-2003 the match is less convincing. The Rat Island event is notable in the history of tsunami forecasting and the DART[®] array. Based on data from early elements of the DART[®] array of the Aleutian Islands, and without the conveniences of the SIFT system for “inversion”, an estimate for the likely impact on the Hawaiian Islands (Titov et al. 2005) demonstrated the utility of direct sea level observation in tsunami forecasting.

The next set of Pacific basin historical events, depicted in Figure 23, are those between Tonga-2006 and Peru-2007. Excluded from the set is the Kuril event of November 15, 2006, that was examined earlier. That event, which was observed at several DART[®] sites, has become a classic for the NCTR modeling group. Tonga-2006 is reported in the NGDC database as producing a 27cm run-up at Crescent City but unfortunately only 6-minute sampled tide gage data are available at Arena Cove. Though the arrival time and first wave shape correspond reasonably well, the amplitude of the observations is considerably less than the model predicts. For Kuril-2006 the tsunami-capable instrument, with its 1-minute sampling, was in place and the early waves of the event were well represented by the forecast model. The same is true of Kuril-2007 which also played an important role in the development of the SIFT forecast tool. Unlike most preceding events whose source mechanism is a reverse thrust fault sending a leading peak toward the offshore DART[®] sites, this was a “normal” thrust event from which a leading trough propagated. As seen in Figure 23, this observed time series at Arena Cove is well matched by the model.

For the Solomon-2007 event the observations were weak and intermittent, though the amplitude of the model signal and its inclusion of larger late waves seems consonant with the data. Also shown in Figure 23, the Peru-2007 event was only weakly felt at Arena Cove and one might be tempted to view the observations as noise. If however the model result is shifted to the right by about ten minutes there is suggestion that the early event history is mimicked. Waves traveling from South America to the U.S. west coast occasionally arrive later than the propagation model predicts, perhaps due to the model bathymetry being smoother than the real ocean. Tsunami waves travel slower in shallower water and in consequence real waves may be delayed in passing through rugged ocean regions such as the Galapagos. Similar delays have been encountered in other forecasts and it remains to be seen whether, as more accurate bathymetric data become available, arrival time forecasts will improve. It should be emphasized that as a percentage of the overall travel time these delays are quite minor.

Late in the same year another event, Chile-2007, occurred off South America. The Arena Cove response, shown in Figure 24, was quite weak and difficult to match with model prediction. Two events from early 2009 are available for study. The predicted signal from the first, near Bird’s Head in Papua-New Guinea, arrived at a noisy period at the Arena Cove tide gage and little if anything can be gleaned from the comparison. Two weeks

later an event of similar magnitude occurred off the Kuril Islands where tsunami waves impacting the west coast frequently originate. As seen in the lower right panel of Figure 24, the Arena Cove FM is reasonably successful in representing that response. Discussion of the Vanuatu 2009 is to be found below.

We now arrive at the events that are most recent at the time this report was written. The Samoa-2009 event was the most damaging to U.S. territory in recent years and, although the DART* array performed well in providing data to provide an accurate forecast, the proximity of the source to U.S. and Western Samoa did not permit any lead time there. This was a sizeable event and, even as far away as Arena Cove, a run-up of 44cm was reported. As seen in Figure 25 the Arena Cove FM performs very well in replicating the early waves, though the later waves may be underestimated. This is an instance in which the reference model (see Figure 18) may, in light of the substantial lead-time inherent in such remote source situations, be a worthwhile option. With a workstation-level computer, the run-time of the FM is presently about 10 hours of clock time for 8 hours of simulation. With advances in computing power or the migration of operational computing to supercomputers it may be possible to depart from the current standard of about 10 minutes per 4-hours of simulation with a FM, perhaps even to run the basin-wide solution in real time with enhanced resolution. Just days after the Samoa-2009 event, another occurred off Vanuatu. Though much less damaging, this event had a new feature to exercise the tsunami community. Vanuatu-2009 was a composite event with two earthquakes in a 15-minute period. Though not yet part of the standard set of historic events for FM evaluation, the separate source characterizations have been established at NCTR (Wei, personal communication). Blending the two forcing histories with an appropriate time delay provides the input needed for a FM (or RM) and the result for the mild response at Arena Cove is quite gratifying. The final historical event analyzed for the first draft of this report is associated with the major earthquake that struck Chile on February 27, 2010. Causing major damage and loss of life locally the tsunami waves propagated widely throughout the Pacific. The waves, seen at DART* 32412, provided a good estimate of the remote hazard, particularly to Hawaii, indicating that wide scale evacuation was not necessary. On the U.S. west coast, noticeable tsunami effects were observed matching predictions. At Arena Cove, had this model been then available and included in the SIFT system, it would have been another point of success for the forecast system in the emergency response to Chile-2010. As seen in Figure 25, there is very close agreement between the FM hind cast and tide gage observation. During the internal NCTR review of this report the Honshu region of Japan was struck, on March 11, 2011, by a huge earthquake, generating a tsunami that caused local devastation and serious impacts throughout the Pacific basin. The FM for Arena Cove was employed in real-time and the results are described briefly in subsection 4.6.

To summarize the analysis of historical events, given above and in subsection 4.6, it would appear that the Arena Cove FM is capable of producing accurate forecasts for this open coast site on the U.S. west coast. Though the observed waves may be difficult to observe accurately at the tide gage during winter storms, the objective of producing credible forecasts of tsunami impact appears to have been met.

4.5 The Mendocino Earthquake of April 25, 1992

Of special interest to northern California is the Mendocino earthquake of April 25, 1992. This has the distinction of being the most recent substantial thrust event on the Cascadia subduction zone. While strike-slip events are commonplace offshore in this region, as shown in Figure 26, it is thrust faults that have the potential to generate significant vertical displacements of the sea floor that cause large tsunamis. The epicenter of this event was on land to the southeast of the plate triple-junction off Cape Mendocino. Uplift of the order of a meter of a 25km stretch of the nearshore, between Cape Mendocino and Punta Gorda to the south was evident in a die-off of intertidal organisms as reported by Carver et al., (1994). Presumably extending offshore too, this deformation is not well represented well by either of the southernmost unit sources (ACSZ-A/B65) now available in the propagation database. The model predictions based on either of these unit sources with an appropriate scale factor for the magnitude 7.2 event underestimate the tide gage signal at Arena Cove, as seen in Figure 26. Another feature of interest of this event, described by González et al. (1995) is that its proximity to shore may have generated a train of coastal-trapped edge waves. Traveling slower than normal tsunami waves taking a deep-water route, the edge waves may have extended the duration of the event at nearby locations to the north and south. This possibility, and the suggestion that the ACSZ source line ought to be extended at least one unit further south, make this an event worth further study. The reference and forecast models for Arena Cove and others existing or planned for the west coast (Eureka, Crescent City, etc.) have a major role in ongoing risk assessment studies for Cascadia.

4.6 The Honshu Tsunami of March 11, 2011

During the NCTR internal review of this report, the severe earthquake and consequent tsunami occurred off the east coast of Honshu, Japan. The SIFT forecast system, ingesting timely data from nearby DART® sites, performed well and provided the basis for appropriate response at those sites for which forecast models were available. Among these was Arena Cove, CA and it seems appropriate to add Honshu-2011 to the suite of historical events for which observations, and both forecast model and reference model results, are available.

The results are shown in Figure 27 where, in the upper panel, the RM and FM time series at the Arena Cove tide gage (drawn as black and red lines respectively) are compared with the 1-minute tide gage record (in blue.) While the largest tsunami waves fortuitously arrived near low water for the U.S. West Coast, and the NCTR models employ mean high water (MHW) in order to represent “worst case” conditions, the agreement is excellent. As at other sites, there was a slight discrepancy in the arrival time (9 minutes in the case of Arena Cove) that has been compensated for in Figure 27. This error is less than 1.6% of the overall travel time and is believed to be associated with the relatively coarse grid of the propagation database which provides the boundary conditions of the finer scale nested FM and RM grids.

After the first few waves the timing and amplitude of the crests and troughs lose synchronicity, both between the RM and FM and between these and the observations. Nonetheless the character of the response is well replicated and the maximum runup agrees well with the reported 1.55m provided by the NGDC database. The latter is the difference between actual and predicted sea level and suggests that, in the case of Arena Cove itself, the forecast wave height is not overly sensitive to the state of the tide, though the extent of inundation may be overstated.

The second row of Figure 27 contrasts the RM and FM solutions at a time, indicated by the green line in the upper panel, where the solutions have begun to diverge. It illustrates that both wave amplitude and tsunami-induced currents are in good agreement through most of the region shown. As before there is some discrepancy near shore at the northern and southern limits of the FM C-grid and near the complex topography off Point Arena Light. The lower panels of the figure contrast the RM and FM predictions for maximum wave amplitude. No reports of amplitude or inundation are available for comparison with these predictions but the agreement would appear to be best near shore for this event. In particular the maxima predicted for Arena Cove and the inundation near the mouth of the Garcia River and much of Manchester Beach match well. If there is error, the FM appears to err on the conservative side, overstating the likely impact.

4.7 Simulation of the remaining Synthetic Mega-events

We conclude this section with a summary of other model runs that were made in order to verify its stability, but which provide useful information on the exposure of Arena Cove to potentially hazardous future events within the Pacific. As noted earlier, the sparse instrumental record of actual events needs to be augmented with credible scenarios to permit risk assessment. While not pretending to be a full-blown risk assessment for the Arena Cove - Manchester Beach area, the full set of mega-tsunamis modeled during stability testing can provide some early estimates.

Results for the set of 19 mega-tsunamis, based on the FM are presented in Figure 28. At the center of each source zone (1000x100km in extent, with the long axis aligned with the local plate boundary and a uniform slip distribution corresponding to an event magnitude of 9.3) a color-coded square represents the impact at Arena Cove. The measure of impact employed is the maximum amplitude of the predicted time series at the reference point (for the Arena Cove FM, the tide gage location near the head of the pier.) There is not any simple relationship between source orientation, location, or great circle distance to Arena Cove; focusing associated with seafloor features can more than compensate for the decay associated with geometric spreading. In Figure 29, FM prediction of the inundation that might result from some of these scenarios are drawn together with (in the lower right panel) an ensemble representing the selection employed in the CalEMA study, whose inundation line is drawn in red.

5.0 Conclusions

In conclusion, good agreement between observations and model predictions for a subset of historical events, including the recent Honshu-2011 tsunami, has been established and the stability of the model for numerous synthetic events has been demonstrated. In particular the reliability of the forecast model, designed to run rapidly in a real time emergency conditions, has been proven by the favorable comparison with reference model predictions, particularly during the early hours of an event. The model will be included in the SIFT system employed operationally at the Tsunami Warning Centers, and will permit the Point Arena – Manchester area to be added to the coastal communities for which forecast capability is available. Additionally it provides a tool of use in risk assessment for the Arena Cove area.

In addition to the scenarios run by the author, and reported here, further tests have been made by other members of the group at NCTR, and will continue to be made by staff at the Warning Centers and others, perhaps in training situations. Among the many related tools developed at NCTR is ComMIT (Community Model Interface for Tsunami, nctr.pmel.noaa.gov/ComMIT/), which provides a highly intuitive graphical environment in which to exercise and explore forecast models for any combination of propagation database unit sources. Were any of these avenues to reveal a problem with the model, its origin (most likely in some quirk of the bathymetric files) would be located and corrected then the revised version re-installed for operational use. The development of the forecast system will be a dynamic process, with new models added (and old ones revisited) from the current list of U.S interests and globally. In the coming years it is expected that further capabilities (for example landslides) will be added as algorithms and methodologies mature.

6. Acknowledgments

Many members of the NCTR GROUP provided valuable assistance in the production of this report. In particular Nicolas Arcos edited the first draft for content and style. CalEMA and other California entities distribute GIS online dataset used in the graphics. The modeling could not proceed without the detailed DEM produced at NGDC by the painstaking combination of numerous bathymetric and topographic surveys. Imagery used in the earlier figures has been reproduced with permission from the California Coastal Records Project (www.californiacoastline.org) and the Mendocino County Historical Society. This publication is [partially] funded by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) under NOAA Cooperative Agreements NA17RJ1232 and NA10OAR4320148, Contribution No. XXXX. It is PMEL Contribution No. NNNN

7. References

- Arcas, D. and B. Uslu (2009): PMEL Tsunami Forecast Series: Vol. 2. A Tsunami Forecast Model for Crescent City, California
- Barberopoulou, A., J. C. Borrero, B. Uslu, M. R. Legg and C. E. Synolakis (2011): A Second Generation of Tsunami Inundation Maps for the State of California. *Pure and Appl Geophys.*
- Barberopoulou, A., J. C. Borrero, B. Uslu, N. Kalogeris, J. D. Goltz, R. I. Wilson, and C.E. Synolakis (2009): New Maps of California to Improve Tsunami Preparedness. *Eos, Trans. AGU, Vol.90, No.16, 21 April 2009*.
See: www.conservation.ca.gov/cgs/geologic_hazards/Tsunami/Inundation_Maps/
- Carver, G.A., A.S. Jayko, D.W. Valentine and W.H. Li (1994): Coastal uplift associated with the 1992 Cape Mendocino earthquake, northern California. *Geology* 22(3), 195-198.
- Driscoll, N.W., Weissel, J.K., and Goff, J.A., 2000, Potential for large-scale submarine slope failure and tsunami generation along the US mid-Atlantic Coast (pdf). *Geology* 28, (5), 407-410.
- Dudley, W.C. and Min Lee (1998): *Tsunami!* University of Hawai'i Press, 362pp.
- Dunbar, P. (2007): Increasing public awareness of natural hazards via the Internet. *Nat. Hazards* 42(3) 529-536, DOI: 10.1007/s11069-006-9072-3
- Fine, I.V., A.B. Rabinovich, B.D. Bornhold, R.E. Thomson, and E.A. Kulikov (2005): The Grand Banks landslide-generated tsunami of November 18, 1929: preliminary analysis and numerical modeling. *Marine Geology* 215, 45–57.
- Friday, D.Z., L.A. Taylor, B.W. Eakins, R.R. Warnken, K.S. Carignan, R.J. Caldwell, E. Lim, and P.R. Medley: Digital Elevation Model of Arena Cove, California: Procedures, Data Sources and Analysis.
www.ngdc.noaa.gov/mgg/inundation/tsunami/data/arena_cove_ca/arena_cove_ca.pdf
- Gica, E., M. Spillane, V.V. Titov, C.D. Chamberlin, and J.C. Newman (2008): Development of the forecast propagation database for NOAA's Short-term Inundation Forecast for Tsunamis (SIFT). NOAA Tech. Memo. OAR PMEL-139, NTIS: PB2008-109391, 89 pp.
- González, F.I., K. Satake, E.F. Boss and H.O. Mofjeld (1995): Edge wave and non-trapped modes of the 25 April 1992 Cape Mendocino tsunami. *Pure and Appl. Geophys.* 144, 409-426, DOI: 10.1007/BF00874375
- Haugan, J. (2005): Dog Holes And Wire Chutes. In: *Maritime Life and Traditions*, No.

29, Winter 2005.

- Lander, J.F., and P.A. Lockridge (1989): United States Tsunamis, 1690 to 1988, Publ. 41–2. National Geophysical Data Center, Boulder, CO, 265 pp.
- O'Brien, M.P. (1946): Preliminary Report on Seismic Sea Waves from Aleutian Earthquake of April 1, 1946, Tech. Rep. HE 116207, Wave Project, Fluid Mechanics Lab., U. Cal. (Berkeley).
- Percival, D.B., D. Arcas, D.W. Denbo, M.C. Eble, E. Gica, H.O. Mofjeld, M.C. Spillane, L. Tang, and V.V. Titov (2009): Extracting tsunami source parameters via inversion of DART® buoy data. NOAA Tech. Memo. OAR PMEL-144, 22 pp.
- Spillane, M.C., E. Gica, V.V. Titov, and H.O. Mofjeld (2008): Tsunameter network design for the U.S. DART® arrays in the Pacific and Atlantic Oceans. NOAA Tech. Memo. OAR PMEL-143, 165 pp.
- Tang, L., C. Chamberlin, E. Tolkova, M. Spillane, V.V. Titov, E.N. Bernard, and H.O. Mofjeld (2006): Assessment of potential tsunami impact for Pearl Harbor, Hawaii. NOAA Tech. Memo. OAR PMEL-131, NTIS: PB2007-100617, 36 pp.
- Tang, L., V.V. Titov, and C.D. Chamberlin (2009): Development, testing, and applications of site-specific tsunami inundation models for real-time forecasting. *J. Geophys. Res.*, 114, C12025, doi: 10.1029/2009JC005476.
- ten Brink, U.S., Lee, H.J., Geist, E.L., Twichell, D.C., (2009): Assessment of tsunami hazard to the U.S. East Coast using relationships between submarine landslides and earthquakes. *Mar. Geol.* 264, 65–73
- Titov, V., and F.I. González (1997): Implementation and testing of the Method of Splitting Tsunami (MOST) model. NOAA Tech. Memo. ERL PMEL-112, NTIS: PB98-122773, NOAA/Pacific Marine Environmental Laboratory, Seattle, WA, 11 pp.
- Titov, V.V., and C.E. Synolakis (1998): Numerical modeling of tidal wave run-up. *J. Waterw. Port Coast. Ocean Eng.*, 124(4), 157–171.
- Titov, V.V., F.I. González, E.N. Bernard, M.C. Eble, H.O. Mofjeld, J.C. Newman, and A.J. Venturato (2005): Real-time tsunami forecasting: Challenges and solutions. *Nat. Hazards*, 35(1), Special Issue, U.S. National Tsunami Hazard Mitigation Program, 41–58.
- U.S. Secretary of War (1914): Harbor of Refuge at Point Arena, or Elsewhere on the Pacific Coast, Between San Francisco and Humboldt Bay, California. Report to 63rd Congress, House of Representatives, Washington D.C., Document No. 1369, 39pp.

Wei, Y., E. Bernard, L. Tang, R. Weiss, V. Titov, C. Moore, M. Spillane, M. Hopkins, and U. Kânoğlu (2008): Real-time experimental forecast of the Peruvian tsunami of August 2007 for U.S. coastlines. *Geophys. Res. Lett.*, 35, L04609, doi: 10.1029/2007GL032250.

DRAFT

FIGURES

DRAFT



Figure 1. The Point Arena area of southern Mendocino County, CA.




Figure 2. Views of present-day Arena Cove and its appearance in the early 1900's.




Figure 3. Extract from the oblique 3-D view of the Arena Cove DEM, provided by NGDC; sites of potential inundation identified by CalEMA are highlighted in red.




Figure 4. Distribution of the historical tsunami sources employed for the development of the Arena Cove forecast model. Those highlighted in red are more extensively investigated using the reference model.




Figure 5. A sample interval from the Arena Cove tsunami-capable tide gage.




Figure 6. The setting of the Arena Cove model.




Figure 7. Nested grid representation for the Reference Model (RM). Rectangles are drawn, when appropriate, to indicate the placement of the next nesting level.




Figure 8. Nested grid representation for the Forecast Model (FM). Rectangles are drawn, when appropriate, to indicate the placement of the next nesting level.




Figure 9. Illustration of the appearance of model instability (upper panel.) The RM C-grid bathymetry in this case required minor modifications, following which the “null” test gave satisfactory results.




Figure 10. Locations of synthetic tsunami scenarios employed in model development.




Figure 11. Comparison of Reference (RM) and Forecast (FM) model results for the ACSZ 56-65 synthetic mega-event, representing the Cascadia Subduction Zone. Time series at the tide gage location in Arena Cove are shown in the upper panel. The lower panels contrast the RM and FM amplitude and velocity fields at the time indicated. Insets are enlargements of Arena Cove.




Figure 12. As in Figure 11, but for the KISZ 01-10 scenario representing Kamchatka.




Figure 13. As in Figure 11, but for the NTSZ 30-30 scenario representing Samoa.




Figure 14. Comparison of the Reference and Forecast Model response to a synthetic moderate event at NTSZ B36 near Samoa.




Figure 15. Comparison of the RM and FM response for the historical Unimak-1946 tsunami (prior to Arena Cove tide gage).




Figure 16. Comparison of the RM and FM response for the historical Alaska-1964 tsunamis (prior to Arena Cove tide gage).




Figure 17. Comparison of RM and FM hind casts of the Kuril 2006 event to sea level fluctuations in Arena Cove, observed by the tsunami-capable tide gage.




Figure 18. Comparison of RM and FM hind casts of the Samoa 2009 event to sea level fluctuations in Arena Cove.




Figure 19. Comparison of RM and FM hind casts of the Chile 2010 event to sea level fluctuations in Arena Cove.




Figure 20. Simulated model response (FM) to the historical Sanriku 1896, Kamchatka 1952, Andreanof 1957 and Chile 1960 events. Proxy observations (in red) are provided where available from other California locations. The Sanriku 1896 result is not expected to match the observed run-up at nearby Mendocino. The state of the tide is shown and times are given as UTC.




Figure 21. Model response (FM) to the East Kuri1 1994, Chile 1995, Kuril 1995 and Irian Jaya 1996 events. Tide gage data (in red) where available is from Arena Cove.




Figure 22. Model response (FM) to the Andreanof 1996, Peru 2001, Hokkaido 2003, and Rat Island 2003 events. Tide gage data (in red) are from Arena Cove.




Figure 23. Model response (FM) to the Tonga 2006, Kuril 2007, Solomons 2007, and Peru 2007 events. Tide gage data (in red) are from Arena Cove




Figure 24. Model response (FM) to the Chile 2007, Papua-New Guinea 2009, Solomons 2009, and Vanuatu 2009 events. Tide gage data (in red) are from Arena Cove.




Figure 25. Model response (FM) to the Samoa 2009 and Chile 2010 events. Tide gage data (in red) are from Arena Cove.




Figure 26. Seismicity in the vicinity of Cape Mendocino and the source mechanisms of notable recent earthquakes, adapted from USGC/NEIC products. The lower panel shows the poor agreement between the model and Arena Cove tide gage observations that is likely due to inadequate source representation in the model.




Figure 27. Comparison of FM real-time forecast and RM hindcast of the Honshu 2011 event with sea level observations in Arena Cove.




Figure 28. Predicted maximum sea level (from the FM) at the Arena Cove tide gauge for the "mega-event" scenarios described in Table 6. Great circle routes are shown in red (with distances in km), and black arrows indicate the normal to the strike direction.




Figure 29. Comparison of RM and FM predictions for inundation of the Arena Cove / Manchester Beach region for selected mega-tsunami scenarios, and (lower right) for the ensemble employed in the CalEMA study.

Appendix A

A1. Reference Model Input (*.in) File for Arena Cove, CA

The following table contains the parameter and file choices used in the input file for the SIFT implementation (most3_facts_nc.in) of the reference model (RM) for Arena Cove, CA.

Parameter/File*	Purpose
0.0010	Minimum amp. of input offshore wave (m)
1.0	Minimum depth of offshore (m)
0.1	Dry land depth of inundation (m)
0.0009	Friction coefficient (n^2)
1	Let A-Grid and B-Grid run up
900.0	Max eta before blow-up (m)
0.6	Time step (sec)
48000	Total number of time steps in run
5	Time steps between A-Grid computations
1	Time steps between B-Grid computations
50	Time steps between output steps
0	Time steps before saving first output step
1	Save output every n-th grid point
ArenaCoveCA_RM_A.most	A-grid bathymetry file
ArenaCoveCA_RM_B.most	B-grid bathymetry file
ArenaCoveCA_RM_C.most	C-grid bathymetry file
./	Directory of source files
./	Directory for output files

* The column headings are not part of most3_facts_nc.in

A2. Forecast Model Input (*.in) File for Arena Cove, CA

The following table contains the parameter and file choices used in the input file for the SIFT implementation (most3_facts_nc.in) of the optimized forecast model (FM) for Arena Cove, CA. When run on an Intel® Xeon® E5670 2.93GHz processor the model simulates four hours in 10.73 minutes, close to the 10-minute target for this metric.

Parameter/File*	Purpose
0.0010	Minimum amp. of input offshore wave (m)
1.0	Minimum depth of offshore (m)
0.1	Dry land depth of inundation (m)
0.0009	Friction coefficient (n^2)
1	Let A-Grid and B-Grid run up
900.0	Max eta before blow-up (m)
1.5	Time step (sec)
19200	Total number of time steps in run
4	Time steps between A-Grid computations
1	Time steps between B-Grid computations
20	Time steps between output steps
0	Time steps before saving first output step
1	Save output every n-th grid point
ArenaCoveCA_FM_A.most	A-grid bathymetry file
ArenaCoveCA_FM_B.most	B-grid bathymetry file
ArenaCoveCA_FM_C.most	C-grid bathymetry file
./	Directory of source files
./	Directory for output files

* The column headings are not part of most3_facts_nc.in

Appendix B Propagation Database: Pacific Ocean Unit Sources

Appendix C Synthetic Testing Report

Glossary

DRAFT

Appendix B

Propagation Database: Pacific Ocean Unit Sources




Figure B.1: Aleutian–Alaska–Cascadia Subduction Zone unit sources.

Table B.1: Earthquake parameters for Aleutian–Alaska–Cascadia Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
acsz-1a	Aleutian–Alaska–Cascadia	164.7994	55.9606	299	17	19.61
acsz-1b	Aleutian–Alaska–Cascadia	164.4310	55.5849	299	17	5
acsz-2a	Aleutian–Alaska–Cascadia	166.3418	55.4016	310.2	17	19.61
acsz-2b	Aleutian–Alaska–Cascadia	165.8578	55.0734	310.2	17	5
acsz-3a	Aleutian–Alaska–Cascadia	167.2939	54.8919	300.2	23.36	24.82
acsz-3b	Aleutian–Alaska–Cascadia	166.9362	54.5356	300.2	23.36	5
acsz-4a	Aleutian–Alaska–Cascadia	168.7131	54.2852	310.2	38.51	25.33
acsz-4b	Aleutian–Alaska–Cascadia	168.3269	54.0168	310.2	24	5
acsz-5a	Aleutian–Alaska–Cascadia	169.7447	53.7808	302.8	37.02	23.54
acsz-5b	Aleutian–Alaska–Cascadia	169.4185	53.4793	302.8	21.77	5
acsz-6a	Aleutian–Alaska–Cascadia	171.0144	53.3054	303.2	35.31	22.92
acsz-6b	Aleutian–Alaska–Cascadia	170.6813	52.9986	303.2	21	5
acsz-7a	Aleutian–Alaska–Cascadia	172.1500	52.8528	298.2	35.56	20.16
acsz-7b	Aleutian–Alaska–Cascadia	171.8665	52.5307	298.2	17.65	5
acsz-8a	Aleutian–Alaska–Cascadia	173.2726	52.4579	290.8	37.92	20.35
acsz-8b	Aleutian–Alaska–Cascadia	173.0681	52.1266	290.8	17.88	5
acsz-9a	Aleutian–Alaska–Cascadia	174.5866	52.1434	289	39.09	21.05
acsz-9b	Aleutian–Alaska–Cascadia	174.4027	51.8138	289	18.73	5
acsz-10a	Aleutian–Alaska–Cascadia	175.8784	51.8526	286.1	40.51	20.87
acsz-10b	Aleutian–Alaska–Cascadia	175.7265	51.5245	286.1	18.51	5
acsz-11a	Aleutian–Alaska–Cascadia	177.1140	51.6488	280	15	17.94
acsz-11b	Aleutian–Alaska–Cascadia	176.9937	51.2215	280	15	5
acsz-12a	Aleutian–Alaska–Cascadia	178.4500	51.5690	273	15	17.94
acsz-12b	Aleutian–Alaska–Cascadia	178.4130	51.1200	273	15	5
acsz-13a	Aleutian–Alaska–Cascadia	179.8550	51.5340	271	15	17.94
acsz-13b	Aleutian–Alaska–Cascadia	179.8420	51.0850	271	15	5
acsz-14a	Aleutian–Alaska–Cascadia	181.2340	51.5780	267	15	17.94
acsz-14b	Aleutian–Alaska–Cascadia	181.2720	51.1290	267	15	5
acsz-15a	Aleutian–Alaska–Cascadia	182.6380	51.6470	265	15	17.94
acsz-15b	Aleutian–Alaska–Cascadia	182.7000	51.2000	265	15	5
acsz-16a	Aleutian–Alaska–Cascadia	184.0550	51.7250	264	15	17.94
acsz-16b	Aleutian–Alaska–Cascadia	184.1280	51.2780	264	15	5
acsz-17a	Aleutian–Alaska–Cascadia	185.4560	51.8170	262	15	17.94
acsz-17b	Aleutian–Alaska–Cascadia	185.5560	51.3720	262	15	5
acsz-18a	Aleutian–Alaska–Cascadia	186.8680	51.9410	261	15	17.94
acsz-18b	Aleutian–Alaska–Cascadia	186.9810	51.4970	261	15	5
acsz-19a	Aleutian–Alaska–Cascadia	188.2430	52.1280	257	15	17.94
acsz-19b	Aleutian–Alaska–Cascadia	188.4060	51.6900	257	15	5
acsz-20a	Aleutian–Alaska–Cascadia	189.5810	52.3550	251	15	17.94
acsz-20b	Aleutian–Alaska–Cascadia	189.8180	51.9300	251	15	5
acsz-21a	Aleutian–Alaska–Cascadia	190.9570	52.6470	251	15	17.94
acsz-21b	Aleutian–Alaska–Cascadia	191.1960	52.2220	251	15	5
acsz-21z	Aleutian–Alaska–Cascadia	190.7399	53.0443	250.8	15	30.88
acsz-22a	Aleutian–Alaska–Cascadia	192.2940	52.9430	247	15	17.94
acsz-22b	Aleutian–Alaska–Cascadia	192.5820	52.5300	247	15	5
acsz-22z	Aleutian–Alaska–Cascadia	192.0074	53.3347	247.8	15	30.88
acsz-23a	Aleutian–Alaska–Cascadia	193.6270	53.3070	245	15	17.94
acsz-23b	Aleutian–Alaska–Cascadia	193.9410	52.9000	245	15	5
acsz-23z	Aleutian–Alaska–Cascadia	193.2991	53.6768	244.6	15	30.88
acsz-24a	Aleutian–Alaska–Cascadia	194.9740	53.6870	245	15	17.94
acsz-24b	Aleutian–Alaska–Cascadia	195.2910	53.2800	245	15	5
acsz-24y	Aleutian–Alaska–Cascadia	194.3645	54.4604	244.4	15	43.82
acsz-24z	Aleutian–Alaska–Cascadia	194.6793	54.0674	244.6	15	30.88

Continued on next page

Table B.1 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
acsz-25a	Aleutian-Alaska-Cascadia	196.4340	54.0760	250	15	17.94
acsz-25b	Aleutian-Alaska-Cascadia	196.6930	53.6543	250	15	5
acsz-25y	Aleutian-Alaska-Cascadia	195.9009	54.8572	247.9	15	43.82
acsz-25z	Aleutian-Alaska-Cascadia	196.1761	54.4536	248.1	15	30.88
acsz-26a	Aleutian-Alaska-Cascadia	197.8970	54.3600	253	15	17.94
acsz-26b	Aleutian-Alaska-Cascadia	198.1200	53.9300	253	15	5
acsz-26y	Aleutian-Alaska-Cascadia	197.5498	55.1934	253.1	15	43.82
acsz-26z	Aleutian-Alaska-Cascadia	197.7620	54.7770	253.3	15	30.88
acsz-27a	Aleutian-Alaska-Cascadia	199.4340	54.5960	256	15	17.94
acsz-27b	Aleutian-Alaska-Cascadia	199.6200	54.1600	256	15	5
acsz-27x	Aleutian-Alaska-Cascadia	198.9736	55.8631	256.5	15	56.24
acsz-27y	Aleutian-Alaska-Cascadia	199.1454	55.4401	256.6	15	43.82
acsz-27z	Aleutian-Alaska-Cascadia	199.3135	55.0170	256.8	15	30.88
acsz-28a	Aleutian-Alaska-Cascadia	200.8820	54.8300	253	15	17.94
acsz-28b	Aleutian-Alaska-Cascadia	201.1080	54.4000	253	15	5
acsz-28x	Aleutian-Alaska-Cascadia	200.1929	56.0559	252.5	15	56.24
acsz-28y	Aleutian-Alaska-Cascadia	200.4167	55.6406	252.7	15	43.82
acsz-28z	Aleutian-Alaska-Cascadia	200.6360	55.2249	252.9	15	30.88
acsz-29a	Aleutian-Alaska-Cascadia	202.2610	55.1330	247	15	17.94
acsz-29b	Aleutian-Alaska-Cascadia	202.5650	54.7200	247	15	5
acsz-29x	Aleutian-Alaska-Cascadia	201.2606	56.2861	245.7	15	56.24
acsz-29y	Aleutian-Alaska-Cascadia	201.5733	55.8888	246	15	43.82
acsz-29z	Aleutian-Alaska-Cascadia	201.8797	55.4908	246.2	15	30.88
acsz-30a	Aleutian-Alaska-Cascadia	203.6040	55.5090	240	15	17.94
acsz-30b	Aleutian-Alaska-Cascadia	203.9970	55.1200	240	15	5
acsz-30w	Aleutian-Alaska-Cascadia	201.9901	56.9855	239.5	15	69.12
acsz-30x	Aleutian-Alaska-Cascadia	202.3851	56.6094	239.8	15	56.24
acsz-30y	Aleutian-Alaska-Cascadia	202.7724	56.2320	240.2	15	43.82
acsz-30z	Aleutian-Alaska-Cascadia	203.1521	55.8534	240.5	15	30.88
acsz-31a	Aleutian-Alaska-Cascadia	204.8950	55.9700	236	15	17.94
acsz-31b	Aleutian-Alaska-Cascadia	205.3400	55.5980	236	15	5
acsz-31w	Aleutian-Alaska-Cascadia	203.0825	57.3740	234.5	15	69.12
acsz-31x	Aleutian-Alaska-Cascadia	203.5408	57.0182	234.9	15	56.24
acsz-31y	Aleutian-Alaska-Cascadia	203.9904	56.6607	235.3	15	43.82
acsz-31z	Aleutian-Alaska-Cascadia	204.4315	56.3016	235.7	15	30.88
acsz-32a	Aleutian-Alaska-Cascadia	206.2080	56.4730	236	15	17.94
acsz-32b	Aleutian-Alaska-Cascadia	206.6580	56.1000	236	15	5
acsz-32w	Aleutian-Alaska-Cascadia	204.4129	57.8908	234.3	15	69.12
acsz-32x	Aleutian-Alaska-Cascadia	204.8802	57.5358	234.7	15	56.24
acsz-32y	Aleutian-Alaska-Cascadia	205.3385	57.1792	235.1	15	43.82
acsz-32z	Aleutian-Alaska-Cascadia	205.7880	56.8210	235.5	15	30.88
acsz-33a	Aleutian-Alaska-Cascadia	207.5370	56.9750	236	15	17.94
acsz-33b	Aleutian-Alaska-Cascadia	207.9930	56.6030	236	15	5
acsz-33w	Aleutian-Alaska-Cascadia	205.7126	58.3917	234.2	15	69.12
acsz-33x	Aleutian-Alaska-Cascadia	206.1873	58.0371	234.6	15	56.24
acsz-33y	Aleutian-Alaska-Cascadia	206.6527	57.6808	235	15	43.82
acsz-33z	Aleutian-Alaska-Cascadia	207.1091	57.3227	235.4	15	30.88
acsz-34a	Aleutian-Alaska-Cascadia	208.9371	57.5124	236	15	17.94
acsz-34b	Aleutian-Alaska-Cascadia	209.4000	57.1400	236	15	5
acsz-34w	Aleutian-Alaska-Cascadia	206.9772	58.8804	233.5	15	69.12
acsz-34x	Aleutian-Alaska-Cascadia	207.4677	58.5291	233.9	15	56.24
acsz-34y	Aleutian-Alaska-Cascadia	207.9485	58.1760	234.3	15	43.82
acsz-34z	Aleutian-Alaska-Cascadia	208.4198	57.8213	234.7	15	30.88
acsz-35a	Aleutian-Alaska-Cascadia	210.2597	58.0441	230	15	17.94
acsz-35b	Aleutian-Alaska-Cascadia	210.8000	57.7000	230	15	5

Continued on next page

Table B.1 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
acsz-35w	Aleutian-Alaska-Cascadia	208.0204	59.3199	228.8	15	69.12
acsz-35x	Aleutian-Alaska-Cascadia	208.5715	58.9906	229.3	15	56.24
acsz-35y	Aleutian-Alaska-Cascadia	209.1122	58.6590	229.7	15	43.82
acsz-35z	Aleutian-Alaska-Cascadia	209.6425	58.3252	230.2	15	30.88
acsz-36a	Aleutian-Alaska-Cascadia	211.3249	58.6565	218	15	17.94
acsz-36b	Aleutian-Alaska-Cascadia	212.0000	58.3800	218	15	5
acsz-36w	Aleutian-Alaska-Cascadia	208.5003	59.5894	215.6	15	69.12
acsz-36x	Aleutian-Alaska-Cascadia	209.1909	59.3342	216.2	15	56.24
acsz-36y	Aleutian-Alaska-Cascadia	209.8711	59.0753	216.8	15	43.82
acsz-36z	Aleutian-Alaska-Cascadia	210.5412	58.8129	217.3	15	30.88
acsz-37a	Aleutian-Alaska-Cascadia	212.2505	59.2720	213.7	15	17.94
acsz-37b	Aleutian-Alaska-Cascadia	212.9519	59.0312	213.7	15	5
acsz-37x	Aleutian-Alaska-Cascadia	210.1726	60.0644	213	15	56.24
acsz-37y	Aleutian-Alaska-Cascadia	210.8955	59.8251	213.7	15	43.82
acsz-37z	Aleutian-Alaska-Cascadia	211.6079	59.5820	214.3	15	30.88
acsz-38a	Aleutian-Alaska-Cascadia	214.6555	60.1351	260.1	0	15
acsz-38b	Aleutian-Alaska-Cascadia	214.8088	59.6927	260.1	0	15
acsz-38y	Aleutian-Alaska-Cascadia	214.3737	60.9838	259	0	15
acsz-38z	Aleutian-Alaska-Cascadia	214.5362	60.5429	259	0	15
acsz-39a	Aleutian-Alaska-Cascadia	216.5607	60.2480	267	0	15
acsz-39b	Aleutian-Alaska-Cascadia	216.6068	59.7994	267	0	15
acsz-40a	Aleutian-Alaska-Cascadia	219.3069	59.7574	310.9	0	15
acsz-40b	Aleutian-Alaska-Cascadia	218.7288	59.4180	310.9	0	15
acsz-41a	Aleutian-Alaska-Cascadia	220.4832	59.3390	300.7	0	15
acsz-41b	Aleutian-Alaska-Cascadia	220.0382	58.9529	300.7	0	15
acsz-42a	Aleutian-Alaska-Cascadia	221.8835	58.9310	298.9	0	15
acsz-42b	Aleutian-Alaska-Cascadia	221.4671	58.5379	298.9	0	15
acsz-43a	Aleutian-Alaska-Cascadia	222.9711	58.6934	282.3	0	15
acsz-43b	Aleutian-Alaska-Cascadia	222.7887	58.2546	282.3	0	15
acsz-44a	Aleutian-Alaska-Cascadia	224.9379	57.9054	340.9	12	11.09
acsz-44b	Aleutian-Alaska-Cascadia	224.1596	57.7617	340.9	7	5
acsz-45a	Aleutian-Alaska-Cascadia	225.4994	57.1634	334.1	12	11.09
acsz-45b	Aleutian-Alaska-Cascadia	224.7740	56.9718	334.1	7	5
acsz-46a	Aleutian-Alaska-Cascadia	226.1459	56.3552	334.1	12	11.09
acsz-46b	Aleutian-Alaska-Cascadia	225.4358	56.1636	334.1	7	5
acsz-47a	Aleutian-Alaska-Cascadia	226.7731	55.5830	332.3	12	11.09
acsz-47b	Aleutian-Alaska-Cascadia	226.0887	55.3785	332.3	7	5
acsz-48a	Aleutian-Alaska-Cascadia	227.4799	54.6763	339.4	12	11.09
acsz-48b	Aleutian-Alaska-Cascadia	226.7713	54.5217	339.4	7	5
acsz-49a	Aleutian-Alaska-Cascadia	227.9482	53.8155	341.2	12	11.09
acsz-49b	Aleutian-Alaska-Cascadia	227.2462	53.6737	341.2	7	5
acsz-50a	Aleutian-Alaska-Cascadia	228.3970	53.2509	324.5	12	11.09
acsz-50b	Aleutian-Alaska-Cascadia	227.8027	52.9958	324.5	7	5
acsz-51a	Aleutian-Alaska-Cascadia	229.1844	52.6297	318.4	12	11.09
acsz-51b	Aleutian-Alaska-Cascadia	228.6470	52.3378	318.4	7	5
acsz-52a	Aleutian-Alaska-Cascadia	230.0306	52.0768	310.9	12	11.09
acsz-52b	Aleutian-Alaska-Cascadia	229.5665	51.7445	310.9	7	5
acsz-53a	Aleutian-Alaska-Cascadia	231.1735	51.5258	310.9	12	11.09
acsz-53b	Aleutian-Alaska-Cascadia	230.7150	51.1935	310.9	7	5
acsz-54a	Aleutian-Alaska-Cascadia	232.2453	50.8809	314.1	12	11.09
acsz-54b	Aleutian-Alaska-Cascadia	231.7639	50.5655	314.1	7	5
acsz-55a	Aleutian-Alaska-Cascadia	233.3066	49.9032	333.7	12	11.09
acsz-55b	Aleutian-Alaska-Cascadia	232.6975	49.7086	333.7	7	5
acsz-56a	Aleutian-Alaska-Cascadia	234.0588	49.1702	315	11	12.82
acsz-56b	Aleutian-Alaska-Cascadia	233.5849	48.8584	315	9	5

Continued on next page

Table B.1 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
acsz-57a	Aleutian-Alaska-Cascadia	234.9041	48.2596	341	11	12.82
acsz-57b	Aleutian-Alaska-Cascadia	234.2797	48.1161	341	9	5
acsz-58a	Aleutian-Alaska-Cascadia	235.3021	47.3812	344	11	12.82
acsz-58b	Aleutian-Alaska-Cascadia	234.6776	47.2597	344	9	5
acsz-59a	Aleutian-Alaska-Cascadia	235.6432	46.5082	345	11	12.82
acsz-59b	Aleutian-Alaska-Cascadia	235.0257	46.3941	345	9	5
acsz-60a	Aleutian-Alaska-Cascadia	235.8640	45.5429	356	11	12.82
acsz-60b	Aleutian-Alaska-Cascadia	235.2363	45.5121	356	9	5
acsz-61a	Aleutian-Alaska-Cascadia	235.9106	44.6227	359	11	12.82
acsz-61b	Aleutian-Alaska-Cascadia	235.2913	44.6150	359	9	5
acsz-62a	Aleutian-Alaska-Cascadia	235.9229	43.7245	359	11	12.82
acsz-62b	Aleutian-Alaska-Cascadia	235.3130	43.7168	359	9	5
acsz-63a	Aleutian-Alaska-Cascadia	236.0220	42.9020	350	11	12.82
acsz-63b	Aleutian-Alaska-Cascadia	235.4300	42.8254	350	9	5
acsz-64a	Aleutian-Alaska-Cascadia	235.9638	41.9818	345	11	12.82
acsz-64b	Aleutian-Alaska-Cascadia	235.3919	41.8677	345	9	5
acsz-65a	Aleutian-Alaska-Cascadia	236.2643	41.1141	345	11	12.82
acsz-65b	Aleutian-Alaska-Cascadia	235.7000	41.0000	345	9	5
acsz-238a	Aleutian-Alaska-Cascadia	213.2878	59.8406	236.8	15	17.94
acsz-238y	Aleutian-Alaska-Cascadia	212.3424	60.5664	236.8	15	43.82
acsz-238z	Aleutian-Alaska-Cascadia	212.8119	60.2035	236.8	15	30.88




Figure B.2: Central and South America Subduction Zone unit sources.

Table B.2: Earthquake parameters for Central and South America Subduction
Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-1a	Central and South America	254.4573	20.8170	359	19	15.4
cssz-1b	Central and South America	254.0035	20.8094	359	12	5
cssz-1z	Central and South America	254.7664	20.8222	359	50	31.67
cssz-2a	Central and South America	254.5765	20.2806	336.8	19	15.4
cssz-2b	Central and South America	254.1607	20.1130	336.8	12	5
cssz-3a	Central and South America	254.8789	19.8923	310.6	18.31	15.27
cssz-3b	Central and South America	254.5841	19.5685	310.6	11.85	5
cssz-4a	Central and South America	255.6167	19.2649	313.4	17.62	15.12
cssz-4b	Central and South America	255.3056	18.9537	313.4	11.68	5
cssz-5a	Central and South America	256.2240	18.8148	302.7	16.92	15
cssz-5b	Central and South America	255.9790	18.4532	302.7	11.54	5
cssz-6a	Central and South America	256.9425	18.4383	295.1	16.23	14.87
cssz-6b	Central and South America	256.7495	18.0479	295.1	11.38	5
cssz-7a	Central and South America	257.8137	18.0339	296.9	15.54	14.74
cssz-7b	Central and South America	257.6079	17.6480	296.9	11.23	5
cssz-8a	Central and South America	258.5779	17.7151	290.4	14.85	14.61
cssz-8b	Central and South America	258.4191	17.3082	290.4	11.08	5
cssz-9a	Central and South America	259.4578	17.4024	290.5	14.15	14.47
cssz-9b	Central and South America	259.2983	16.9944	290.5	10.92	5
cssz-10a	Central and South America	260.3385	17.0861	290.8	13.46	14.34
cssz-10b	Central and South America	260.1768	16.6776	290.8	10.77	5
cssz-11a	Central and South America	261.2255	16.7554	291.8	12.77	14.21
cssz-11b	Central and South America	261.0556	16.3487	291.8	10.62	5
cssz-12a	Central and South America	262.0561	16.4603	288.9	12.08	14.08
cssz-12b	Central and South America	261.9082	16.0447	288.9	10.46	5
cssz-13a	Central and South America	262.8638	16.2381	283.2	11.38	13.95
cssz-13b	Central and South America	262.7593	15.8094	283.2	10.31	5
cssz-14a	Central and South America	263.6066	16.1435	272.1	10.69	13.81
cssz-14b	Central and South America	263.5901	15.7024	272.1	10.15	5
cssz-15a	Central and South America	264.8259	15.8829	293	10	13.68
cssz-15b	Central and South America	264.6462	15.4758	293	10	5
cssz-15y	Central and South America	265.1865	16.6971	293	10	31.05
cssz-15z	Central and South America	265.0060	16.2900	293	10	22.36
cssz-16a	Central and South America	265.7928	15.3507	304.9	15	15.82
cssz-16b	Central and South America	265.5353	14.9951	304.9	12.5	5
cssz-16y	Central and South America	266.3092	16.0619	304.9	15	41.7
cssz-16z	Central and South America	266.0508	15.7063	304.9	15	28.76
cssz-17a	Central and South America	266.4947	14.9019	299.5	20	17.94
cssz-17b	Central and South America	266.2797	14.5346	299.5	15	5
cssz-17y	Central and South America	266.9259	15.6365	299.5	20	52.14
cssz-17z	Central and South America	266.7101	15.2692	299.5	20	35.04
cssz-18a	Central and South America	267.2827	14.4768	298	21.5	17.94
cssz-18b	Central and South America	267.0802	14.1078	298	15	5
cssz-18y	Central and South America	267.6888	15.2148	298	21.5	54.59
cssz-18z	Central and South America	267.4856	14.8458	298	21.5	36.27
cssz-19a	Central and South America	268.0919	14.0560	297.6	23	17.94
cssz-19b	Central and South America	267.8943	13.6897	297.6	15	5
cssz-19y	Central and South America	268.4880	14.7886	297.6	23	57.01
cssz-19z	Central and South America	268.2898	14.4223	297.6	23	37.48
cssz-20a	Central and South America	268.8929	13.6558	296.2	24	17.94
cssz-20b	Central and South America	268.7064	13.2877	296.2	15	5
cssz-20y	Central and South America	269.1796	14.2206	296.2	45.5	73.94
cssz-20z	Central and South America	269.0362	13.9382	296.2	45.5	38.28

Continued on next page

Table B.2 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-21a	Central and South America	269.6797	13.3031	292.6	25	17.94
cssz-21b	Central and South America	269.5187	12.9274	292.6	15	5
cssz-21x	Central and South America	269.8797	13.7690	292.6	68	131.8
cssz-21y	Central and South America	269.8130	13.6137	292.6	68	85.43
cssz-21z	Central and South America	269.7463	13.4584	292.6	68	39.07
cssz-22a	Central and South America	270.4823	13.0079	288.6	25	17.94
cssz-22b	Central and South America	270.3492	12.6221	288.6	15	5
cssz-22x	Central and South America	270.6476	13.4864	288.6	68	131.8
cssz-22y	Central and South America	270.5925	13.3269	288.6	68	85.43
cssz-22z	Central and South America	270.5374	13.1674	288.6	68	39.07
cssz-23a	Central and South America	271.3961	12.6734	292.4	25	17.94
cssz-23b	Central and South America	271.2369	12.2972	292.4	15	5
cssz-23x	Central and South America	271.5938	13.1399	292.4	68	131.8
cssz-23y	Central and South America	271.5279	12.9844	292.4	68	85.43
cssz-23z	Central and South America	271.4620	12.8289	292.4	68	39.07
cssz-24a	Central and South America	272.3203	12.2251	300.2	25	17.94
cssz-24b	Central and South America	272.1107	11.8734	300.2	15	5
cssz-24x	Central and South America	272.5917	12.6799	300.2	67	131.1
cssz-24y	Central and South America	272.5012	12.5283	300.2	67	85.1
cssz-24z	Central and South America	272.4107	12.3767	300.2	67	39.07
cssz-25a	Central and South America	273.2075	11.5684	313.8	25	17.94
cssz-25b	Central and South America	272.9200	11.2746	313.8	15	5
cssz-25x	Central and South America	273.5950	11.9641	313.8	66	130.4
cssz-25y	Central and South America	273.4658	11.8322	313.8	66	84.75
cssz-25z	Central and South America	273.3366	11.7003	313.8	66	39.07
cssz-26a	Central and South America	273.8943	10.8402	320.4	25	17.94
cssz-26b	Central and South America	273.5750	10.5808	320.4	15	5
cssz-26x	Central and South America	274.3246	11.1894	320.4	66	130.4
cssz-26y	Central and South America	274.1811	11.0730	320.4	66	84.75
cssz-26z	Central and South America	274.0377	10.9566	320.4	66	39.07
cssz-27a	Central and South America	274.4569	10.2177	316.1	25	17.94
cssz-27b	Central and South America	274.1590	9.9354	316.1	15	5
cssz-27z	Central and South America	274.5907	10.3444	316.1	66	39.07
cssz-28a	Central and South America	274.9586	9.8695	297.1	22	14.54
cssz-28b	Central and South America	274.7661	9.4988	297.1	11	5
cssz-28z	Central and South America	275.1118	10.1643	297.1	42.5	33.27
cssz-29a	Central and South America	275.7686	9.4789	296.6	19	11.09
cssz-29b	Central and South America	275.5759	9.0992	296.6	7	5
cssz-30a	Central and South America	276.6346	8.9973	302.2	19	9.36
cssz-30b	Central and South America	276.4053	8.6381	302.2	5	5
cssz-31a	Central and South America	277.4554	8.4152	309.1	19	7.62
cssz-31b	Central and South America	277.1851	8.0854	309.1	3	5
cssz-31z	Central and South America	277.7260	8.7450	309.1	19	23.9
cssz-32a	Central and South America	278.1112	7.9425	303	18.67	8.49
cssz-32b	Central and South America	277.8775	7.5855	303	4	5
cssz-32z	Central and South America	278.3407	8.2927	303	21.67	24.49
cssz-33a	Central and South America	278.7082	7.6620	287.6	18.33	10.23
cssz-33b	Central and South America	278.5785	7.2555	287.6	6	5
cssz-33z	Central and South America	278.8328	8.0522	287.6	24.33	25.95
cssz-34a	Central and South America	279.3184	7.5592	269.5	18	17.94
cssz-34b	Central and South America	279.3223	7.1320	269.5	15	5
cssz-35a	Central and South America	280.0039	7.6543	255.9	17.67	14.54
cssz-35b	Central and South America	280.1090	7.2392	255.9	11	5
cssz-35x	Central and South America	279.7156	8.7898	255.9	29.67	79.22
cssz-35y	Central and South America	279.8118	8.4113	255.9	29.67	54.47

Continued on next page

Table B.2 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-35z	Central and South America	279.9079	8.0328	255.9	29.67	29.72
cssz-36a	Central and South America	281.2882	7.6778	282.5	17.33	11.09
cssz-36b	Central and South America	281.1948	7.2592	282.5	7	5
cssz-36x	Central and South America	281.5368	8.7896	282.5	32.33	79.47
cssz-36y	Central and South America	281.4539	8.4190	282.5	32.33	52.73
cssz-36z	Central and South America	281.3710	8.0484	282.5	32.33	25.99
cssz-37a	Central and South America	282.5252	6.8289	326.9	17	10.23
cssz-37b	Central and South America	282.1629	6.5944	326.9	6	5
cssz-38a	Central and South America	282.9469	5.5973	355.4	17	10.23
cssz-38b	Central and South America	282.5167	5.5626	355.4	6	5
cssz-39a	Central and South America	282.7236	4.3108	24.13	17	10.23
cssz-39b	Central and South America	282.3305	4.4864	24.13	6	5
cssz-39z	Central and South America	283.0603	4.1604	24.13	35	24.85
cssz-40a	Central and South America	282.1940	3.3863	35.28	17	10.23
cssz-40b	Central and South America	281.8427	3.6344	35.28	6	5
cssz-40y	Central and South America	282.7956	2.9613	35.28	35	53.52
cssz-40z	Central and South America	282.4948	3.1738	35.28	35	24.85
cssz-41a	Central and South America	281.6890	2.6611	34.27	17	10.23
cssz-41b	Central and South America	281.3336	2.9030	34.27	6	5
cssz-41z	Central and South America	281.9933	2.4539	34.27	35	24.85
cssz-42a	Central and South America	281.2266	1.9444	31.29	17	10.23
cssz-42b	Central and South America	280.8593	2.1675	31.29	6	5
cssz-42z	Central and South America	281.5411	1.7533	31.29	35	24.85
cssz-43a	Central and South America	280.7297	1.1593	33.3	17	10.23
cssz-43b	Central and South America	280.3706	1.3951	33.3	6	5
cssz-43z	Central and South America	281.0373	0.9573	33.3	35	24.85
cssz-44a	Central and South America	280.3018	0.4491	28.8	17	10.23
cssz-44b	Central and South America	279.9254	0.6560	28.8	6	5
cssz-45a	Central and South America	279.9083	-0.3259	26.91	10	8.49
cssz-45b	Central and South America	279.5139	-0.1257	26.91	4	5
cssz-46a	Central and South America	279.6461	-0.9975	15.76	10	8.49
cssz-46b	Central and South America	279.2203	-0.8774	15.76	4	5
cssz-47a	Central and South America	279.4972	-1.7407	6.9	10	8.49
cssz-47b	Central and South America	279.0579	-1.6876	6.9	4	5
cssz-48a	Central and South America	279.3695	-2.6622	8.96	10	8.49
cssz-48b	Central and South America	278.9321	-2.5933	8.96	4	5
cssz-48y	Central and South America	280.2444	-2.8000	8.96	10	25.85
cssz-48z	Central and South America	279.8070	-2.7311	8.96	10	17.17
cssz-49a	Central and South America	279.1852	-3.6070	13.15	10	8.49
cssz-49b	Central and South America	278.7536	-3.5064	13.15	4	5
cssz-49y	Central and South America	280.0486	-3.8082	13.15	10	25.85
cssz-49z	Central and South America	279.6169	-3.7076	13.15	10	17.17
cssz-50a	Central and South America	279.0652	-4.3635	4.78	10.33	9.64
cssz-50b	Central and South America	278.6235	-4.3267	4.78	5.33	5
cssz-51a	Central and South America	279.0349	-5.1773	359.4	10.67	10.81
cssz-51b	Central and South America	278.5915	-5.1817	359.4	6.67	5
cssz-52a	Central and South America	279.1047	-5.9196	349.8	11	11.96
cssz-52b	Central and South America	278.6685	-5.9981	349.8	8	5
cssz-53a	Central and South America	279.3044	-6.6242	339.2	10.25	11.74
cssz-53b	Central and South America	278.8884	-6.7811	339.2	7.75	5
cssz-53y	Central and South America	280.1024	-6.3232	339.2	19.25	37.12
cssz-53z	Central and South America	279.7035	-6.4737	339.2	19.25	20.64
cssz-54a	Central and South America	279.6256	-7.4907	340.8	9.5	11.53
cssz-54b	Central and South America	279.2036	-7.6365	340.8	7.5	5
cssz-54y	Central and South America	280.4267	-7.2137	340.8	20.5	37.29

Continued on next page

Table B.2 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-54z	Central and South America	280.0262	-7.3522	340.8	20.5	19.78
cssz-55a	Central and South America	279.9348	-8.2452	335.4	8.75	11.74
cssz-55b	Central and South America	279.5269	-8.4301	335.4	7.75	5
cssz-55x	Central and South America	281.0837	-7.7238	335.4	21.75	56.4
cssz-55y	Central and South America	280.7009	-7.8976	335.4	21.75	37.88
cssz-55z	Central and South America	280.3180	-8.0714	335.4	21.75	19.35
cssz-56a	Central and South America	280.3172	-8.9958	331.6	8	11.09
cssz-56b	Central and South America	279.9209	-9.2072	331.6	7	5
cssz-56x	Central and South America	281.4212	-8.4063	331.6	23	57.13
cssz-56y	Central and South America	281.0534	-8.6028	331.6	23	37.59
cssz-56z	Central and South America	280.6854	-8.7993	331.6	23	18.05
cssz-57a	Central and South America	280.7492	-9.7356	328.7	8.6	10.75
cssz-57b	Central and South America	280.3640	-9.9663	328.7	6.6	5
cssz-57x	Central and South America	281.8205	-9.0933	328.7	23.4	57.94
cssz-57y	Central and South America	281.4636	-9.3074	328.7	23.4	38.08
cssz-57z	Central and South America	281.1065	-9.5215	328.7	23.4	18.22
cssz-58a	Central and South America	281.2275	-10.5350	330.5	9.2	10.4
cssz-58b	Central and South America	280.8348	-10.7532	330.5	6.2	5
cssz-58y	Central and South America	281.9548	-10.1306	330.5	23.8	38.57
cssz-58z	Central and South America	281.5913	-10.3328	330.5	23.8	18.39
cssz-59a	Central and South America	281.6735	-11.2430	326.2	9.8	10.05
cssz-59b	Central and South America	281.2982	-11.4890	326.2	5.8	5
cssz-59y	Central and South America	282.3675	-10.7876	326.2	24.2	39.06
cssz-59z	Central and South America	282.0206	-11.0153	326.2	24.2	18.56
cssz-60a	Central and South America	282.1864	-11.9946	326.5	10.4	9.71
cssz-60b	Central and South America	281.8096	-12.2384	326.5	5.4	5
cssz-60y	Central and South America	282.8821	-11.5438	326.5	24.6	39.55
cssz-60z	Central and South America	282.5344	-11.7692	326.5	24.6	18.73
cssz-61a	Central and South America	282.6944	-12.7263	325.5	11	9.36
cssz-61b	Central and South America	282.3218	-12.9762	325.5	5	5
cssz-61y	Central and South America	283.3814	-12.2649	325.5	25	40.03
cssz-61z	Central and South America	283.0381	-12.4956	325.5	25	18.9
cssz-62a	Central and South America	283.1980	-13.3556	319	11	9.79
cssz-62b	Central and South America	282.8560	-13.6451	319	5.5	5
cssz-62y	Central and South America	283.8178	-12.8300	319	27	42.03
cssz-62z	Central and South America	283.5081	-13.0928	319	27	19.33
cssz-63a	Central and South America	283.8032	-14.0147	317.9	11	10.23
cssz-63b	Central and South America	283.4661	-14.3106	317.9	6	5
cssz-63z	Central and South America	284.1032	-13.7511	317.9	29	19.77
cssz-64a	Central and South America	284.4144	-14.6482	315.7	13	11.96
cssz-64b	Central and South America	284.0905	-14.9540	315.7	8	5
cssz-65a	Central and South America	285.0493	-15.2554	313.2	15	13.68
cssz-65b	Central and South America	284.7411	-15.5715	313.2	10	5
cssz-66a	Central and South America	285.6954	-15.7816	307.7	14.5	13.68
cssz-66b	Central and South America	285.4190	-16.1258	307.7	10	5
cssz-67a	Central and South America	286.4127	-16.2781	304.3	14	13.68
cssz-67b	Central and South America	286.1566	-16.6381	304.3	10	5
cssz-67z	Central and South America	286.6552	-15.9365	304.3	23	25.78
cssz-68a	Central and South America	287.2481	-16.9016	311.8	14	13.68
cssz-68b	Central and South America	286.9442	-17.2264	311.8	10	5
cssz-68z	Central and South America	287.5291	-16.6007	311.8	26	25.78
cssz-69a	Central and South America	287.9724	-17.5502	314.9	14	13.68
cssz-69b	Central and South America	287.6496	-17.8590	314.9	10	5
cssz-69y	Central and South America	288.5530	-16.9934	314.9	29	50.02
cssz-69z	Central and South America	288.2629	-17.2718	314.9	29	25.78

Continued on next page

Table B.2 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-70a	Central and South America	288.6731	-18.2747	320.4	14	13.25
cssz-70b	Central and South America	288.3193	-18.5527	320.4	9.5	5
cssz-70y	Central and South America	289.3032	-17.7785	320.4	30	50.35
cssz-70z	Central and South America	288.9884	-18.0266	320.4	30	25.35
cssz-71a	Central and South America	289.3089	-19.1854	333.2	14	12.82
cssz-71b	Central and South America	288.8968	-19.3820	333.2	9	5
cssz-71y	Central and South America	290.0357	-18.8382	333.2	31	50.67
cssz-71z	Central and South America	289.6725	-19.0118	333.2	31	24.92
cssz-72a	Central and South America	289.6857	-20.3117	352.4	14	12.54
cssz-72b	Central and South America	289.2250	-20.3694	352.4	8.67	5
cssz-72z	Central and South America	290.0882	-20.2613	352.4	32	24.63
cssz-73a	Central and South America	289.7731	-21.3061	358.9	14	12.24
cssz-73b	Central and South America	289.3053	-21.3142	358.9	8.33	5
cssz-73z	Central and South America	290.1768	-21.2991	358.9	33	24.34
cssz-74a	Central and South America	289.7610	-22.2671	3.06	14	11.96
cssz-74b	Central and South America	289.2909	-22.2438	3.06	8	5
cssz-75a	Central and South America	289.6982	-23.1903	4.83	14.09	11.96
cssz-75b	Central and South America	289.2261	-23.1536	4.83	8	5
cssz-76a	Central and South America	289.6237	-24.0831	4.67	14.18	11.96
cssz-76b	Central and South America	289.1484	-24.0476	4.67	8	5
cssz-77a	Central and South America	289.5538	-24.9729	4.3	14.27	11.96
cssz-77b	Central and South America	289.0750	-24.9403	4.3	8	5
cssz-78a	Central and South America	289.4904	-25.8621	3.86	14.36	11.96
cssz-78b	Central and South America	289.0081	-25.8328	3.86	8	5
cssz-79a	Central and South America	289.3491	-26.8644	11.34	14.45	11.96
cssz-79b	Central and South America	288.8712	-26.7789	11.34	8	5
cssz-80a	Central and South America	289.1231	-27.7826	14.16	14.54	11.96
cssz-80b	Central and South America	288.6469	-27.6762	14.16	8	5
cssz-81a	Central and South America	288.8943	-28.6409	13.19	14.63	11.96
cssz-81b	Central and South America	288.4124	-28.5417	13.19	8	5
cssz-82a	Central and South America	288.7113	-29.4680	9.68	14.72	11.96
cssz-82b	Central and South America	288.2196	-29.3950	9.68	8	5
cssz-83a	Central and South America	288.5944	-30.2923	5.36	14.81	11.96
cssz-83b	Central and South America	288.0938	-30.2517	5.36	8	5
cssz-84a	Central and South America	288.5223	-31.1639	3.8	14.9	11.96
cssz-84b	Central and South America	288.0163	-31.1351	3.8	8	5
cssz-85a	Central and South America	288.4748	-32.0416	2.55	15	11.96
cssz-85b	Central and South America	287.9635	-32.0223	2.55	8	5
cssz-86a	Central and South America	288.3901	-33.0041	7.01	15	11.96
cssz-86b	Central and South America	287.8768	-32.9512	7.01	8	5
cssz-87a	Central and South America	288.1050	-34.0583	19.4	15	11.96
cssz-87b	Central and South America	287.6115	-33.9142	19.4	8	5
cssz-88a	Central and South America	287.5309	-35.0437	32.81	15	11.96
cssz-88b	Central and South America	287.0862	-34.8086	32.81	8	5
cssz-88z	Central and South America	287.9308	-35.2545	32.81	30	24.9
cssz-89a	Central and South America	287.2380	-35.5993	14.52	16.67	11.96
cssz-89b	Central and South America	286.7261	-35.4914	14.52	8	5
cssz-89z	Central and South America	287.7014	-35.6968	14.52	30	26.3
cssz-90a	Central and South America	286.8442	-36.5645	22.64	18.33	11.96
cssz-90b	Central and South America	286.3548	-36.4004	22.64	8	5
cssz-90z	Central and South America	287.2916	-36.7142	22.64	30	27.68
cssz-91a	Central and South America	286.5925	-37.2488	10.9	20	11.96
cssz-91b	Central and South America	286.0721	-37.1690	10.9	8	5
cssz-91z	Central and South America	287.0726	-37.3224	10.9	30	29.06
cssz-92a	Central and South America	286.4254	-38.0945	8.23	20	11.96

Continued on next page

Table B.2 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-92b	Central and South America	285.8948	-38.0341	8.23	8	5
cssz-92z	Central and South America	286.9303	-38.1520	8.23	26.67	29.06
cssz-93a	Central and South America	286.2047	-39.0535	13.46	20	11.96
cssz-93b	Central and South America	285.6765	-38.9553	13.46	8	5
cssz-93z	Central and South America	286.7216	-39.1495	13.46	23.33	29.06
cssz-94a	Central and South America	286.0772	-39.7883	3.4	20	11.96
cssz-94b	Central and South America	285.5290	-39.7633	3.4	8	5
cssz-94z	Central and South America	286.6255	-39.8133	3.4	20	29.06
cssz-95a	Central and South America	285.9426	-40.7760	9.84	20	11.96
cssz-95b	Central and South America	285.3937	-40.7039	9.84	8	5
cssz-95z	Central and South America	286.4921	-40.8481	9.84	20	29.06
cssz-96a	Central and South America	285.7839	-41.6303	7.6	20	11.96
cssz-96b	Central and South America	285.2245	-41.5745	7.6	8	5
cssz-96x	Central and South America	287.4652	-41.7977	7.6	20	63.26
cssz-96y	Central and South America	286.9043	-41.7419	7.6	20	46.16
cssz-96z	Central and South America	286.3439	-41.6861	7.6	20	29.06
cssz-97a	Central and South America	285.6695	-42.4882	5.3	20	11.96
cssz-97b	Central and South America	285.0998	-42.4492	5.3	8	5
cssz-97x	Central and South America	287.3809	-42.6052	5.3	20	63.26
cssz-97y	Central and South America	286.8101	-42.5662	5.3	20	46.16
cssz-97z	Central and South America	286.2396	-42.5272	5.3	20	29.06
cssz-98a	Central and South America	285.5035	-43.4553	10.53	20	11.96
cssz-98b	Central and South America	284.9322	-43.3782	10.53	8	5
cssz-98x	Central and South America	287.2218	-43.6866	10.53	20	63.26
cssz-98y	Central and South America	286.6483	-43.6095	10.53	20	46.16
cssz-98z	Central and South America	286.0755	-43.5324	10.53	20	29.06
cssz-99a	Central and South America	285.3700	-44.2595	4.86	20	11.96
cssz-99b	Central and South America	284.7830	-44.2237	4.86	8	5
cssz-99x	Central and South America	287.1332	-44.3669	4.86	20	63.26
cssz-99y	Central and South America	286.5451	-44.3311	4.86	20	46.16
cssz-99z	Central and South America	285.9574	-44.2953	4.86	20	29.06
cssz-100a	Central and South America	285.2713	-45.1664	5.68	20	11.96
cssz-100b	Central and South America	284.6758	-45.1246	5.68	8	5
cssz-100x	Central and South America	287.0603	-45.2918	5.68	20	63.26
cssz-100y	Central and South America	286.4635	-45.2500	5.68	20	46.16
cssz-100z	Central and South America	285.8672	-45.2082	5.68	20	29.06
cssz-101a	Central and South America	285.3080	-45.8607	352.6	20	9.36
cssz-101b	Central and South America	284.7067	-45.9152	352.6	5	5
cssz-101y	Central and South America	286.5089	-45.7517	352.6	20	43.56
cssz-101z	Central and South America	285.9088	-45.8062	352.6	20	26.46
cssz-102a	Central and South America	285.2028	-47.1185	17.72	5	9.36
cssz-102b	Central and South America	284.5772	-46.9823	17.72	5	5
cssz-102y	Central and South America	286.4588	-47.3909	17.72	5	18.07
cssz-102z	Central and South America	285.8300	-47.2547	17.72	5	13.72
cssz-103a	Central and South America	284.7075	-48.0396	23.37	7.5	11.53
cssz-103b	Central and South America	284.0972	-47.8630	23.37	7.5	5
cssz-103x	Central and South America	286.5511	-48.5694	23.37	7.5	31.11
cssz-103y	Central and South America	285.9344	-48.3928	23.37	7.5	24.58
cssz-103z	Central and South America	285.3199	-48.2162	23.37	7.5	18.05
cssz-104a	Central and South America	284.3440	-48.7597	14.87	10	13.68
cssz-104b	Central and South America	283.6962	-48.6462	14.87	10	5
cssz-104x	Central and South America	286.2962	-49.1002	14.87	10	39.73
cssz-104y	Central and South America	285.6440	-48.9867	14.87	10	31.05
cssz-104z	Central and South America	284.9933	-48.8732	14.87	10	22.36
cssz-105a	Central and South America	284.2312	-49.4198	0.25	9.67	13.4

Continued on next page

Table B.2 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
cssz-105b	Central and South America	283.5518	-49.4179	0.25	9.67	5
cssz-105x	Central and South America	286.2718	-49.4255	0.25	9.67	38.59
cssz-105y	Central and South America	285.5908	-49.4236	0.25	9.67	30.2
cssz-105z	Central and South America	284.9114	-49.4217	0.25	9.67	21.8
cssz-106a	Central and South America	284.3730	-50.1117	347.5	9.25	13.04
cssz-106b	Central and South America	283.6974	-50.2077	347.5	9.25	5
cssz-106x	Central and South America	286.3916	-49.8238	347.5	9.25	37.15
cssz-106y	Central and South America	285.7201	-49.9198	347.5	9.25	29.11
cssz-106z	Central and South America	285.0472	-50.0157	347.5	9.25	21.07
cssz-107a	Central and South America	284.7130	-50.9714	346.5	9	12.82
cssz-107b	Central and South America	284.0273	-51.0751	346.5	9	5
cssz-107x	Central and South America	286.7611	-50.6603	346.5	9	36.29
cssz-107y	Central and South America	286.0799	-50.7640	346.5	9	28.47
cssz-107z	Central and South America	285.3972	-50.8677	346.5	9	20.64
cssz-108a	Central and South America	285.0378	-51.9370	352	8.67	12.54
cssz-108b	Central and South America	284.3241	-51.9987	352	8.67	5
cssz-108x	Central and South America	287.1729	-51.7519	352	8.67	35.15
cssz-108y	Central and South America	286.4622	-51.8136	352	8.67	27.61
cssz-108z	Central and South America	285.7505	-51.8753	352	8.67	20.07
cssz-109a	Central and South America	285.2635	-52.8439	353.1	8.33	12.24
cssz-109b	Central and South America	284.5326	-52.8974	353.1	8.33	5
cssz-109x	Central and South America	287.4508	-52.6834	353.1	8.33	33.97
cssz-109y	Central and South America	286.7226	-52.7369	353.1	8.33	26.73
cssz-109z	Central and South America	285.9935	-52.7904	353.1	8.33	19.49
cssz-110a	Central and South America	285.5705	-53.4139	334.2	8	11.96
cssz-110b	Central and South America	284.8972	-53.6076	334.2	8	5
cssz-110x	Central and South America	287.5724	-52.8328	334.2	8	32.83
cssz-110y	Central and South America	286.9081	-53.0265	334.2	8	25.88
cssz-110z	Central and South America	286.2408	-53.2202	334.2	8	18.92
cssz-111a	Central and South America	286.1627	-53.8749	313.8	8	11.96
cssz-111b	Central and South America	285.6382	-54.1958	313.8	8	5
cssz-111x	Central and South America	287.7124	-52.9122	313.8	8	32.83
cssz-111y	Central and South America	287.1997	-53.2331	313.8	8	25.88
cssz-111z	Central and South America	286.6832	-53.5540	313.8	8	18.92
cssz-112a	Central and South America	287.3287	-54.5394	316.4	8	11.96
cssz-112b	Central and South America	286.7715	-54.8462	316.4	8	5
cssz-112x	Central and South America	288.9756	-53.6190	316.4	8	32.83
cssz-112y	Central and South America	288.4307	-53.9258	316.4	8	25.88
cssz-112z	Central and South America	287.8817	-54.2326	316.4	8	18.92
cssz-113a	Central and South America	288.3409	-55.0480	307.6	8	11.96
cssz-113b	Central and South America	287.8647	-55.4002	307.6	8	5
cssz-113x	Central and South America	289.7450	-53.9914	307.6	8	32.83
cssz-113y	Central and South America	289.2810	-54.3436	307.6	8	25.88
cssz-113z	Central and South America	288.8130	-54.6958	307.6	8	18.92
cssz-114a	Central and South America	289.5342	-55.5026	301.5	8	11.96
cssz-114b	Central and South America	289.1221	-55.8819	301.5	8	5
cssz-114x	Central and South America	290.7472	-54.3647	301.5	8	32.83
cssz-114y	Central and South America	290.3467	-54.7440	301.5	8	25.88
cssz-114z	Central and South America	289.9424	-55.1233	301.5	8	18.92
cssz-115a	Central and South America	290.7682	-55.8485	292.7	8	11.96
cssz-115b	Central and South America	290.4608	-56.2588	292.7	8	5
cssz-115x	Central and South America	291.6714	-54.6176	292.7	8	32.83
cssz-115y	Central and South America	291.3734	-55.0279	292.7	8	25.88
cssz-115z	Central and South America	291.0724	-55.4382	292.7	8	18.92




Figure B.3: Eastern Philippines Subduction Zone unit sources.

Table B.3: Earthquake parameters for Eastern Philippines Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
epsz-1a	Eastern Philippines	128.5521	2.3289	153.6	44.2	27.62
epsz-1b	Eastern Philippines	128.8408	2.4720	153.6	26.9	5
epsz-2a	Eastern Philippines	128.1943	3.1508	151.9	45.9	32.44
epsz-2b	Eastern Philippines	128.4706	3.2979	151.9	32.8	5.35
epsz-3a	Eastern Philippines	127.8899	4.0428	155.2	57.3	40.22
epsz-3b	Eastern Philippines	128.1108	4.1445	155.2	42.7	6.31
epsz-4a	Eastern Philippines	127.6120	4.8371	146.8	71.4	48.25
epsz-4b	Eastern Philippines	127.7324	4.9155	146.8	54.8	7.39
epsz-5a	Eastern Philippines	127.3173	5.7040	162.9	79.9	57.4
epsz-5b	Eastern Philippines	127.3930	5.7272	162.9	79.4	8.25
epsz-6a	Eastern Philippines	126.6488	6.6027	178.9	48.6	45.09
epsz-6b	Eastern Philippines	126.9478	6.6085	178.9	48.6	7.58
epsz-7a	Eastern Philippines	126.6578	7.4711	175.8	50.7	45.52
epsz-7b	Eastern Philippines	126.9439	7.4921	175.8	50.7	6.83
epsz-8a	Eastern Philippines	126.6227	8.2456	163.3	56.7	45.6
epsz-8b	Eastern Philippines	126.8614	8.3164	163.3	48.9	7.92
epsz-9a	Eastern Philippines	126.2751	9.0961	164.1	47	43.59
epsz-9b	Eastern Philippines	126.5735	9.1801	164.1	44.9	8.3
epsz-10a	Eastern Philippines	125.9798	9.9559	164.5	43.1	42.25
epsz-10b	Eastern Philippines	126.3007	10.0438	164.5	43.1	8.09
epsz-11a	Eastern Philippines	125.6079	10.6557	155	37.8	38.29
epsz-11b	Eastern Philippines	125.9353	10.8059	155	37.8	7.64
epsz-12a	Eastern Philippines	125.4697	11.7452	172.1	36	37.01
epsz-12b	Eastern Philippines	125.8374	11.7949	172.1	36	7.62
epsz-13a	Eastern Philippines	125.2238	12.1670	141.5	32.4	33.87
epsz-13b	Eastern Philippines	125.5278	12.4029	141.5	32.4	7.08
epsz-14a	Eastern Philippines	124.6476	13.1365	158.2	23	25.92
epsz-14b	Eastern Philippines	125.0421	13.2898	158.2	23	6.38
epsz-15a	Eastern Philippines	124.3107	13.9453	156.1	24.1	26.51
epsz-15b	Eastern Philippines	124.6973	14.1113	156.1	24.1	6.09
epsz-16a	Eastern Philippines	123.8998	14.4025	140.3	19.5	21.69
epsz-16b	Eastern Philippines	124.2366	14.6728	140.3	19.5	5
epsz-17a	Eastern Philippines	123.4604	14.7222	117.6	15.3	18.19
epsz-17b	Eastern Philippines	123.6682	15.1062	117.6	15.3	5
epsz-18a	Eastern Philippines	123.3946	14.7462	67.4	15	17.94
epsz-18b	Eastern Philippines	123.2219	15.1467	67.4	15	5
epsz-19a	Eastern Philippines	121.3638	15.7400	189.6	15	17.94
epsz-19b	Eastern Philippines	121.8082	15.6674	189.6	15	5
epsz-20a	Eastern Philippines	121.6833	16.7930	203.3	15	17.94
epsz-20b	Eastern Philippines	122.0994	16.6216	203.3	15	5
epsz-21a	Eastern Philippines	121.8279	17.3742	184.2	15	17.94
epsz-21b	Eastern Philippines	122.2814	17.3425	184.2	15	5




Figure B.4: Kamchatka-Kuril-Japan-Izu-Mariana-Yap Subduction Zone unit sources.

Table B.4: Earthquake parameters for Kamchatka-Kuril-Japan-Izu-Mariana-Yap Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
kisz-1a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	162.4318	55.5017	195	29	26.13
kisz-1b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	163.1000	55.4000	195	25	5
kisz-1y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	161.0884	55.7050	195	29	74.61
kisz-1z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	161.7610	55.6033	195	29	50.37
kisz-2a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	161.9883	54.6784	200	29	26.13
kisz-2b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	162.6247	54.5440	200	25	5
kisz-2y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.7072	54.9471	200	29	74.61
kisz-2z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	161.3488	54.8127	200	29	50.37
kisz-3a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	161.4385	53.8714	204	29	26.13
kisz-3b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	162.0449	53.7116	204	25	5
kisz-3y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.2164	54.1910	204	29	74.61
kisz-3z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.8286	54.0312	204	29	50.37
kisz-4a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.7926	53.1087	210	29	26.13
kisz-4b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	161.3568	52.9123	210	25	5
kisz-4y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	159.6539	53.5015	210	29	74.61
kisz-4z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.2246	53.3051	210	29	50.37
kisz-5a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.0211	52.4113	218	29	26.13
kisz-5b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	160.5258	52.1694	218	25	5
kisz-5y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	159.0005	52.8950	218	29	74.61
kisz-5z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	159.5122	52.6531	218	29	50.37
kisz-6a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	159.1272	51.7034	218	29	26.13
kisz-6b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	159.6241	51.4615	218	25	5
kisz-6y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	158.1228	52.1871	218	29	74.61
kisz-6z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	158.6263	51.9452	218	29	50.37
kisz-7a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	158.2625	50.9549	214	29	26.13
kisz-7b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	158.7771	50.7352	214	25	5
kisz-7y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	157.2236	51.3942	214	29	74.61
kisz-7z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	157.7443	51.1745	214	29	50.37
kisz-8a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	157.4712	50.2459	218	31	27.7
kisz-8b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	157.9433	50.0089	218	27	5
kisz-8y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	156.5176	50.7199	218	31	79.2
kisz-8z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	156.9956	50.4829	218	31	53.45
kisz-9a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	156.6114	49.5583	220	31	27.7
kisz-9b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	157.0638	49.3109	220	27	5
kisz-9y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	155.6974	50.0533	220	31	79.2
kisz-9z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	156.1556	49.8058	220	31	53.45
kisz-10a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	155.7294	48.8804	221	31	27.7
kisz-10b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	156.1690	48.6278	221	27	5
kisz-10y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	154.8413	49.3856	221	31	79.2
kisz-10z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	155.2865	49.1330	221	31	53.45
kisz-11a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	154.8489	48.1821	219	31	27.7
kisz-11b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	155.2955	47.9398	219	27	5
kisz-11y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.9472	48.6667	219	31	79.2
kisz-11z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	154.3991	48.4244	219	31	53.45
kisz-11c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	156.0358	47.5374	39	57.89	4.602
kisz-12a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.9994	47.4729	217	31	27.7
kisz-12b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	154.4701	47.2320	217	27	5
kisz-12y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.0856	47.9363	217	31	79.2
kisz-12z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.5435	47.7046	217	31	53.45
kisz-12c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	155.2208	46.8473	37	57.89	4.602
kisz-13a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.2239	46.7564	218	31	27.7
kisz-13b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.6648	46.5194	218	27	5

Continued on next page

Table B.4 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
kisz-13y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	152.3343	47.2304	218	31	79.2
kisz-13z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	152.7801	46.9934	218	31	53.45
kisz-13c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	154.3957	46.1257	38	57.89	4.602
kisz-14a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	152.3657	46.1514	225	23	24.54
kisz-14b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	152.7855	45.8591	225	23	5
kisz-14y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.5172	46.7362	225	23	63.62
kisz-14z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.9426	46.4438	225	23	44.08
kisz-14c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.4468	45.3976	45	57.89	4.602
kisz-15a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.4663	45.5963	233	25	23.73
kisz-15b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.8144	45.2712	233	22	5
kisz-15y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.7619	46.2465	233	25	65.99
kisz-15z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.1151	45.9214	233	25	44.86
kisz-16a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.4572	45.0977	237	25	23.73
kisz-16b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.7694	44.7563	237	22	5
kisz-16y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.8253	45.7804	237	25	65.99
kisz-16z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.1422	45.4390	237	25	44.86
kisz-17a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.3989	44.6084	237	25	23.73
kisz-17b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.7085	44.2670	237	22	5
kisz-17y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.7723	45.2912	237	25	65.99
kisz-17z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.0865	44.9498	237	25	44.86
kisz-18a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.3454	44.0982	235	25	23.73
kisz-18b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.6687	43.7647	235	22	5
kisz-18y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.6915	44.7651	235	25	65.99
kisz-18z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.0194	44.4316	235	25	44.86
kisz-19a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.3262	43.5619	233	25	23.73
kisz-19b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.6625	43.2368	233	22	5
kisz-19y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.6463	44.2121	233	25	65.99
kisz-19z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.9872	43.8870	233	25	44.86
kisz-20a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.3513	43.0633	237	25	23.73
kisz-20b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.6531	42.7219	237	22	5
kisz-20y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.7410	43.7461	237	25	65.99
kisz-20z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.0470	43.4047	237	25	44.86
kisz-21a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.3331	42.5948	239	25	23.73
kisz-21b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.6163	42.2459	239	22	5
kisz-21y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.7603	43.2927	239	25	65.99
kisz-21z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.0475	42.9438	239	25	44.86
kisz-22a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.3041	42.1631	242	25	23.73
kisz-22b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.5605	41.8037	242	22	5
kisz-22y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.7854	42.8819	242	25	65.99
kisz-22z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.0455	42.5225	242	25	44.86
kisz-23a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.2863	41.3335	202	21	21.28
kisz-23b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.8028	41.1764	202	19	5
kisz-23v	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.6816	42.1189	202	21	110.9
kisz-23w	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.2050	41.9618	202	21	92.95
kisz-23x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.7273	41.8047	202	21	75.04
kisz-23y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.2482	41.6476	202	21	57.12
kisz-23z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7679	41.4905	202	21	39.2
kisz-24a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.9795	40.3490	185	21	21.28
kisz-24b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.5273	40.3125	185	19	5
kisz-24x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.3339	40.4587	185	21	75.04
kisz-24y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.8827	40.4221	185	21	57.12
kisz-24z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.4312	40.3856	185	21	39.2
kisz-25a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.8839	39.4541	185	21	21.28
kisz-25b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.4246	39.4176	185	19	5
kisz-25y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.8012	39.5272	185	21	57.12

Continued on next page

Table B.4 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
kisz-25z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.3426	39.4907	185	21	39.2
kisz-26a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7622	38.5837	188	21	21.28
kisz-26b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.2930	38.5254	188	19	5
kisz-26x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.1667	38.7588	188	21	75.04
kisz-26y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6990	38.7004	188	21	57.12
kisz-26z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.2308	38.6421	188	21	39.2
kisz-27a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.5320	37.7830	198	21	21.28
kisz-27b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.0357	37.6534	198	19	5
kisz-27x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.0142	38.1717	198	21	75.04
kisz-27y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.5210	38.0421	198	21	57.12
kisz-27z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.0269	37.9126	198	21	39.2
kisz-28a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.1315	37.0265	208	21	21.28
kisz-28b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.5941	36.8297	208	19	5
kisz-28x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.7348	37.6171	208	21	75.04
kisz-28y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.2016	37.4202	208	21	57.12
kisz-28z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6671	37.2234	208	21	39.2
kisz-29a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.5970	36.2640	211	21	21.28
kisz-29b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.0416	36.0481	211	19	5
kisz-29y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.7029	36.6960	211	21	57.12
kisz-29z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.1506	36.4800	211	21	39.2
kisz-30a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.0553	35.4332	205	21	21.28
kisz-30b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.5207	35.2560	205	19	5
kisz-30y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.1204	35.7876	205	21	57.12
kisz-30z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.5883	35.6104	205	21	39.2
kisz-31a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.6956	34.4789	190	22	22.1
kisz-31b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.1927	34.4066	190	20	5
kisz-31v	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.2025	34.8405	190	22	115.8
kisz-31w	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.7021	34.7682	190	22	97.02
kisz-31x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	139.2012	34.6958	190	22	78.29
kisz-31y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	139.6997	34.6235	190	22	59.56
kisz-31z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.1979	34.5512	190	22	40.83
kisz-32a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.0551	33.0921	180	32	23.48
kisz-32b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.5098	33.0921	180	21.69	5
kisz-33a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.0924	32.1047	173.8	27.65	20.67
kisz-33b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.5596	32.1473	173.8	18.27	5
kisz-34a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.1869	31.1851	172.1	25	18.26
kisz-34b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6585	31.2408	172.1	15.38	5
kisz-35a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.4154	30.1707	163	25	17.12
kisz-35b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.8662	30.2899	163	14.03	5
kisz-36a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6261	29.2740	161.7	25.73	18.71
kisz-36b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.0670	29.4012	161.7	15.91	5
kisz-37a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.0120	28.3322	154.7	20	14.54
kisz-37b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.4463	28.5124	154.7	11	5
kisz-38a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.2254	27.6946	170.3	20	14.54
kisz-38b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.6955	27.7659	170.3	11	5
kisz-39a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.3085	26.9127	177.2	24.23	17.42
kisz-39b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7674	26.9325	177.2	14.38	5
kisz-40a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.2673	26.1923	189.4	26.49	22.26
kisz-40b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7090	26.1264	189.4	20.2	5
kisz-41a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.1595	25.0729	173.7	22.07	19.08
kisz-41b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.6165	25.1184	173.7	16.36	5
kisz-42a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7641	23.8947	143.5	21.54	18.4
kisz-42b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.1321	24.1432	143.5	15.54	5
kisz-43a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.5281	23.0423	129.2	23.02	18.77
kisz-43b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.8128	23.3626	129.2	15.99	5

Continued on next page

Table B.4 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
kisz-44a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.2230	22.5240	134.6	28.24	18.56
kisz-44b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.5246	22.8056	134.6	15.74	5
kisz-45a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.0895	21.8866	125.8	36.73	22.79
kisz-45b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.3171	22.1785	125.8	20.84	5
kisz-46a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.6972	21.3783	135.9	30.75	20.63
kisz-46b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.9954	21.6469	135.9	18.22	5
kisz-47a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.0406	20.9341	160.1	29.87	19.62
kisz-47b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.4330	21.0669	160.1	17	5
kisz-48a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.3836	20.0690	158	32.75	19.68
kisz-48b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.7567	20.2108	158	17.07	5
kisz-49a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.6689	19.3123	164.5	25.07	21.41
kisz-49b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.0846	19.4212	164.5	19.16	5
kisz-50a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.9297	18.5663	172.1	22	22.1
kisz-50b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.3650	18.6238	172.1	20	5
kisz-51a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.9495	17.7148	175.1	22.06	22.04
kisz-51b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.3850	17.7503	175.1	19.93	5
kisz-52a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.9447	16.8869	180	25.51	18.61
kisz-52b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.3683	16.8869	180	15.79	5
kisz-53a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.8626	16.0669	185.2	27.39	18.41
kisz-53b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.2758	16.0309	185.2	15.56	5
kisz-54a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.7068	15.3883	199.1	28.12	20.91
kisz-54b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.0949	15.2590	199.1	18.56	5
kisz-55a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.4717	14.6025	204.3	29.6	26.27
kisz-55b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.8391	14.4415	204.3	25.18	5
kisz-56a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.1678	13.9485	217.4	32.04	26.79
kisz-56b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.4789	13.7170	217.4	25.84	5
kisz-57a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.6515	13.5576	235.8	37	24.54
kisz-57b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.8586	13.2609	235.8	23	5
kisz-58a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.9648	12.9990	237.8	37.72	24.54
kisz-58b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.1589	12.6984	237.8	23	5
kisz-59a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.1799	12.6914	242.9	34.33	22.31
kisz-59b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.3531	12.3613	242.9	20.25	5
kisz-60a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.3687	12.3280	244.9	30.9	20.62
kisz-60b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.5355	11.9788	244.9	18.2	5
kisz-61a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7051	12.1507	261.8	35.41	25.51
kisz-61b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7582	11.7883	261.8	24.22	5
kisz-62a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6301	11.8447	245.7	39.86	34.35
kisz-62b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.7750	11.5305	245.7	35.94	5
kisz-63a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.8923	11.5740	256.2	42	38.46
kisz-63b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.9735	11.2498	256.2	42	5
kisz-64a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.1387	11.6028	269.6	42.48	38.77
kisz-64b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.1410	11.2716	269.6	42.48	5
kisz-65a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	139.4595	11.5883	288.7	44.16	39.83
kisz-65b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	139.3541	11.2831	288.7	44.16	5
kisz-66a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.1823	11.2648	193.1	45	40.36
kisz-66b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.4977	11.1929	193.1	45	5
kisz-67a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.9923	10.3398	189.8	45	40.36
kisz-67b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.3104	10.2856	189.8	45	5
kisz-68a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.7607	9.6136	201.7	45	40.36
kisz-68b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.0599	9.4963	201.7	45	5
kisz-69a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.4537	8.8996	213.5	45	40.36
kisz-69b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.7215	8.7241	213.5	45	5
kisz-70a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.0191	8.2872	226.5	45	40.36
kisz-70b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.2400	8.0569	226.5	45	5
kisz-71a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	136.3863	7.9078	263.9	45	40.36

Continued on next page

Table B.4 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
kisz-71b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	136.4202	7.5920	263.9	45	5
kisz-72a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	135.6310	7.9130	276.9	45	40.36
kisz-72b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	135.5926	7.5977	276.9	45	5
kisz-73a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	134.3296	7.4541	224	45	40.36
kisz-73b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	134.5600	7.2335	224	45	5
kisz-74a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	133.7125	6.8621	228.1	45	40.36
kisz-74b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	133.9263	6.6258	228.1	45	5
kisz-75a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	133.0224	6.1221	217.7	45	40.36
kisz-75b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	133.2751	5.9280	217.7	45	5




Figure B.5: Manus–Oceanic Convergent Boundary Subduction Zone unit sources.

Table B.5: Earthquake parameters for Manus–Oceanic Convergent Boundary Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
mosz-1a	Manus	154.0737	-4.8960	140.2	15	15.88
mosz-1b	Manus	154.4082	-4.6185	140.2	15	2.94
mosz-2a	Manus	153.5589	-4.1575	140.2	15	15.91
mosz-2b	Manus	153.8931	-3.8800	140.2	15	2.97
mosz-3a	Manus	153.0151	-3.3716	143.9	15	16.64
mosz-3b	Manus	153.3662	-3.1160	143.9	15	3.7
mosz-4a	Manus	152.4667	-3.0241	127.7	15	17.32
mosz-4b	Manus	152.7321	-2.6806	127.7	15	4.38
mosz-5a	Manus	151.8447	-2.7066	114.3	15	17.57
mosz-5b	Manus	152.0235	-2.3112	114.3	15	4.63
mosz-6a	Manus	151.0679	-2.2550	115	15	17.66
mosz-6b	Manus	151.2513	-1.8618	115	15	4.72
mosz-7a	Manus	150.3210	-2.0236	107.2	15	17.73
mosz-7b	Manus	150.4493	-1.6092	107.2	15	4.79
mosz-8a	Manus	149.3226	-1.6666	117.8	15	17.83
mosz-8b	Manus	149.5251	-1.2829	117.8	15	4.89
mosz-9a	Manus	148.5865	-1.3017	112.7	15	17.84
mosz-9b	Manus	148.7540	-0.9015	112.7	15	4.9
mosz-10a	Manus	147.7760	-1.1560	108	15	17.78
mosz-10b	Manus	147.9102	-0.7434	108	15	4.84
mosz-11a	Manus	146.9596	-1.1226	102.5	15	17.54
mosz-11b	Manus	147.0531	-0.6990	102.5	15	4.6
mosz-12a	Manus	146.2858	-1.1820	87.48	15	17.29
mosz-12b	Manus	146.2667	-0.7486	87.48	15	4.35
mosz-13a	Manus	145.4540	-1.3214	83.75	15	17.34
mosz-13b	Manus	145.4068	-0.8901	83.75	15	4.4
mosz-14a	Manus	144.7151	-1.5346	75.09	15	17.21
mosz-14b	Manus	144.6035	-1.1154	75.09	15	4.27
mosz-15a	Manus	143.9394	-1.8278	70.43	15	16.52
mosz-15b	Manus	143.7940	-1.4190	70.43	15	3.58
mosz-16a	Manus	143.4850	-2.2118	50.79	15	15.86
mosz-16b	Manus	143.2106	-1.8756	50.79	15	2.92
mosz-17a	Manus	143.1655	-2.7580	33	15	16.64
mosz-17b	Manus	142.8013	-2.5217	33	15	3.7




Figure B.6: New Guinea Subduction Zone unit sources.

Table B.6: Earthquake parameters for New Guinea Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
ngsz-1a	New Guinea	143.6063	-4.3804	120	29	25.64
ngsz-1b	New Guinea	143.8032	-4.0402	120	29	1.4
ngsz-2a	New Guinea	142.9310	-3.9263	114	27.63	20.1
ngsz-2b	New Guinea	143.0932	-3.5628	114	21.72	1.6
ngsz-3a	New Guinea	142.1076	-3.5632	114	20.06	18.73
ngsz-3b	New Guinea	142.2795	-3.1778	114	15.94	5
ngsz-4a	New Guinea	141.2681	-3.2376	114	21	17.76
ngsz-4b	New Guinea	141.4389	-2.8545	114	14.79	5
ngsz-5a	New Guinea	140.4592	-2.8429	114	21.26	16.14
ngsz-5b	New Guinea	140.6296	-2.4605	114	12.87	5
ngsz-6a	New Guinea	139.6288	-2.4960	114	22.72	15.4
ngsz-6b	New Guinea	139.7974	-2.1175	114	12	5
ngsz-7a	New Guinea	138.8074	-2.1312	114	21.39	15.4
ngsz-7b	New Guinea	138.9776	-1.7491	114	12	5
ngsz-8a	New Guinea	138.0185	-1.7353	113.1	18.79	15.14
ngsz-8b	New Guinea	138.1853	-1.3441	113.1	11.7	5
ngsz-9a	New Guinea	137.1805	-1.5037	111	15.24	13.23
ngsz-9b	New Guinea	137.3358	-1.0991	111	9.47	5
ngsz-10a	New Guinea	136.3418	-1.1774	111	13.51	11.09
ngsz-10b	New Guinea	136.4983	-0.7697	111	7	5
ngsz-11a	New Guinea	135.4984	-0.8641	111	11.38	12.49
ngsz-11b	New Guinea	135.6562	-0.4530	111	8.62	5
ngsz-12a	New Guinea	134.6759	-0.5216	110.5	10	13.68
ngsz-12b	New Guinea	134.8307	-0.1072	110.5	10	5
ngsz-13a	New Guinea	133.3065	-1.0298	99.5	10	13.68
ngsz-13b	New Guinea	133.3795	-0.5935	99.5	10	5
ngsz-14a	New Guinea	132.4048	-0.8816	99.5	10	13.68
ngsz-14b	New Guinea	132.4778	-0.4453	99.5	10	5
ngsz-15a	New Guinea	131.5141	-0.7353	99.5	10	13.68
ngsz-15b	New Guinea	131.5871	-0.2990	99.5	10	5




Figure B.7: New Zealand–Kermadec–Tonga Subduction Zone unit sources.

Table B.7: Earthquake parameters for New Zealand–Kermadec–Tonga Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
ntsz-1a	New Zealand–Tonga	174.0985	-41.3951	258.6	24	25.34
ntsz-1b	New Zealand–Tonga	174.2076	-41.7973	258.6	24	5
ntsz-2a	New Zealand–Tonga	175.3289	-41.2592	260.6	29.38	23.17
ntsz-2b	New Zealand–Tonga	175.4142	-41.6454	260.6	21.31	5
ntsz-3a	New Zealand–Tonga	176.2855	-40.9950	250.7	29.54	21.74
ntsz-3b	New Zealand–Tonga	176.4580	-41.3637	250.7	19.56	5
ntsz-4a	New Zealand–Tonga	177.0023	-40.7679	229.4	24.43	18.87
ntsz-4b	New Zealand–Tonga	177.3552	-41.0785	229.4	16.1	5
ntsz-5a	New Zealand–Tonga	177.4114	-40.2396	210	18.8	19.29
ntsz-5b	New Zealand–Tonga	177.8951	-40.4525	210	16.61	5
ntsz-6a	New Zealand–Tonga	177.8036	-39.6085	196.7	18.17	15.8
ntsz-6b	New Zealand–Tonga	178.3352	-39.7310	196.7	12.48	5
ntsz-7a	New Zealand–Tonga	178.1676	-38.7480	197	28.1	17.85
ntsz-7b	New Zealand–Tonga	178.6541	-38.8640	197	14.89	5
ntsz-8a	New Zealand–Tonga	178.6263	-37.8501	201.4	31.47	18.78
ntsz-8b	New Zealand–Tonga	179.0788	-37.9899	201.4	16	5
ntsz-9a	New Zealand–Tonga	178.9833	-36.9770	202.2	29.58	20.02
ntsz-9b	New Zealand–Tonga	179.4369	-37.1245	202.2	17.48	5
ntsz-10a	New Zealand–Tonga	179.5534	-36.0655	210.6	32.1	20.72
ntsz-10b	New Zealand–Tonga	179.9595	-36.2593	210.6	18.32	5
ntsz-11a	New Zealand–Tonga	179.9267	-35.3538	201.7	25	16.09
ntsz-11b	New Zealand–Tonga	180.3915	-35.5040	201.7	12.81	5
ntsz-12a	New Zealand–Tonga	180.4433	-34.5759	201.2	25	15.46
ntsz-12b	New Zealand–Tonga	180.9051	-34.7230	201.2	12.08	5
ntsz-13a	New Zealand–Tonga	180.7990	-33.7707	199.8	25.87	19.06
ntsz-13b	New Zealand–Tonga	181.2573	-33.9073	199.8	16.33	5
ntsz-14a	New Zealand–Tonga	181.2828	-32.9288	202.4	31.28	22.73
ntsz-14b	New Zealand–Tonga	181.7063	-33.0751	202.4	20.77	5
ntsz-15a	New Zealand–Tonga	181.4918	-32.0035	205.4	32.33	22.64
ntsz-15b	New Zealand–Tonga	181.8967	-32.1665	205.4	20.66	5
ntsz-16a	New Zealand–Tonga	181.9781	-31.2535	205.5	34.29	23.59
ntsz-16b	New Zealand–Tonga	182.3706	-31.4131	205.5	21.83	5
ntsz-17a	New Zealand–Tonga	182.4819	-30.3859	210.3	37.6	25.58
ntsz-17b	New Zealand–Tonga	182.8387	-30.5655	210.3	24.3	5
ntsz-18a	New Zealand–Tonga	182.8176	-29.6545	201.6	37.65	26.13
ntsz-18b	New Zealand–Tonga	183.1985	-29.7856	201.6	25	5
ntsz-19a	New Zealand–Tonga	183.0622	-28.8739	195.7	34.41	26.13
ntsz-19b	New Zealand–Tonga	183.4700	-28.9742	195.7	25	5
ntsz-20a	New Zealand–Tonga	183.2724	-28.0967	188.8	38	26.13
ntsz-20b	New Zealand–Tonga	183.6691	-28.1508	188.8	25	5
ntsz-21a	New Zealand–Tonga	183.5747	-27.1402	197.1	32.29	24.83
ntsz-21b	New Zealand–Tonga	183.9829	-27.2518	197.1	23.37	5
ntsz-22a	New Zealand–Tonga	183.6608	-26.4975	180	29.56	18.63
ntsz-22b	New Zealand–Tonga	184.0974	-26.4975	180	15.82	5
ntsz-23a	New Zealand–Tonga	183.7599	-25.5371	185.8	32.42	20.56
ntsz-23b	New Zealand–Tonga	184.1781	-25.5752	185.8	18.13	5
ntsz-24a	New Zealand–Tonga	183.9139	-24.6201	188.2	33.31	23.73
ntsz-24b	New Zealand–Tonga	184.3228	-24.6734	188.2	22	5
ntsz-25a	New Zealand–Tonga	184.1266	-23.5922	198.5	29.34	19.64
ntsz-25b	New Zealand–Tonga	184.5322	-23.7163	198.5	17.03	5
ntsz-26a	New Zealand–Tonga	184.6613	-22.6460	211.7	30.26	19.43
ntsz-26b	New Zealand–Tonga	185.0196	-22.8497	211.7	16.78	5
ntsz-27a	New Zealand–Tonga	185.0879	-21.9139	207.9	31.73	20.67

Continued on next page

Table B.7 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
ntsz-27b	New Zealand–Tonga	185.4522	-22.0928	207.9	18.27	5
ntsz-28a	New Zealand–Tonga	185.4037	-21.1758	200.5	32.44	21.76
ntsz-28b	New Zealand–Tonga	185.7849	-21.3084	200.5	19.58	5
ntsz-29a	New Zealand–Tonga	185.8087	-20.2629	206.4	32.47	20.4
ntsz-29b	New Zealand–Tonga	186.1710	-20.4312	206.4	17.94	5
ntsz-30a	New Zealand–Tonga	186.1499	-19.5087	200.9	32.98	22.46
ntsz-30b	New Zealand–Tonga	186.5236	-19.6432	200.9	20.44	5
ntsz-31a	New Zealand–Tonga	186.3538	-18.7332	193.9	34.41	21.19
ntsz-31b	New Zealand–Tonga	186.7339	-18.8221	193.9	18.89	5
ntsz-32a	New Zealand–Tonga	186.5949	-17.8587	194.1	30	19.12
ntsz-32b	New Zealand–Tonga	186.9914	-17.9536	194.1	16.4	5
ntsz-33a	New Zealand–Tonga	186.8172	-17.0581	190	33.15	23.34
ntsz-33b	New Zealand–Tonga	187.2047	-17.1237	190	21.52	5
ntsz-34a	New Zealand–Tonga	186.7814	-16.2598	182.1	15	13.41
ntsz-34b	New Zealand–Tonga	187.2330	-16.2759	182.1	9.68	5
ntsz-34c	New Zealand–Tonga	187.9697	-16.4956	7.62	57.06	6.571
ntsz-35a	New Zealand–Tonga	186.8000	-15.8563	149.8	15	12.17
ntsz-35b	New Zealand–Tonga	187.1896	-15.6384	149.8	8.24	5
ntsz-35c	New Zealand–Tonga	187.8776	-15.6325	342.4	57.06	6.571
ntsz-36a	New Zealand–Tonga	186.5406	-15.3862	123.9	40.44	36.72
ntsz-36b	New Zealand–Tonga	186.7381	-15.1025	123.9	39.38	5
ntsz-36c	New Zealand–Tonga	187.3791	-14.9234	307	57.06	6.571
ntsz-37a	New Zealand–Tonga	185.9883	-14.9861	102	68.94	30.99
ntsz-37b	New Zealand–Tonga	186.0229	-14.8282	102	31.32	5
ntsz-38a	New Zealand–Tonga	185.2067	-14.8259	88.4	80	26.13
ntsz-38b	New Zealand–Tonga	185.2044	-14.7479	88.4	25	5
ntsz-39a	New Zealand–Tonga	184.3412	-14.9409	82.55	80	26.13
ntsz-39b	New Zealand–Tonga	184.3307	-14.8636	82.55	25	5




Figure B.8: New Britain–Solomons–Vanuatu Zone unit sources.

Table B.8: Earthquake parameters for New Britain–Solomons–Vanuatu Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
nvsh-1a	New Britain–Vanuatu	148.6217	-6.4616	243.2	32.34	15.69
nvsh-1b	New Britain–Vanuatu	148.7943	-6.8002	234.2	12.34	5
nvsh-2a	New Britain–Vanuatu	149.7218	-6.1459	260.1	35.1	16.36
nvsh-2b	New Britain–Vanuatu	149.7856	-6.5079	260.1	13.13	5
nvsh-3a	New Britain–Vanuatu	150.4075	-5.9659	245.7	42.35	18.59
nvsh-3b	New Britain–Vanuatu	150.5450	-6.2684	245.7	15.77	5
nvsh-4a	New Britain–Vanuatu	151.1095	-5.5820	238.2	42.41	23.63
nvsh-4b	New Britain–Vanuatu	151.2851	-5.8639	238.2	21.88	5
nvsh-5a	New Britain–Vanuatu	152.0205	-5.1305	247.7	49.22	32.39
nvsh-5b	New Britain–Vanuatu	152.1322	-5.4020	247.7	33.22	5
nvsh-6a	New Britain–Vanuatu	153.3450	-5.1558	288.6	53.53	33.59
nvsh-6b	New Britain–Vanuatu	153.2595	-5.4089	288.6	34.87	5
nvsh-7a	New Britain–Vanuatu	154.3814	-5.6308	308.3	39.72	19.18
nvsh-7b	New Britain–Vanuatu	154.1658	-5.9017	308.3	16.48	5
nvsh-8a	New Britain–Vanuatu	155.1097	-6.3511	317.2	45.33	22.92
nvsh-8b	New Britain–Vanuatu	154.8764	-6.5656	317.2	21	5
nvsh-9a	New Britain–Vanuatu	155.5027	-6.7430	290.5	48.75	22.92
nvsh-9b	New Britain–Vanuatu	155.3981	-7.0204	290.5	21	5
nvsh-10a	New Britain–Vanuatu	156.4742	-7.2515	305.9	36.88	27.62
nvsh-10b	New Britain–Vanuatu	156.2619	-7.5427	305.9	26.9	5
nvsh-11a	New Britain–Vanuatu	157.0830	-7.8830	305.4	32.97	29.72
nvsh-11b	New Britain–Vanuatu	156.8627	-8.1903	305.4	29.63	5
nvsh-12a	New Britain–Vanuatu	157.6537	-8.1483	297.9	37.53	28.57
nvsh-12b	New Britain–Vanuatu	157.4850	-8.4630	297.9	28.13	5
nvsh-13a	New Britain–Vanuatu	158.5089	-8.5953	302.7	33.62	23.02
nvsh-13b	New Britain–Vanuatu	158.3042	-8.9099	302.7	21.12	5
nvsh-14a	New Britain–Vanuatu	159.1872	-8.9516	293.3	38.44	34.06
nvsh-14b	New Britain–Vanuatu	159.0461	-9.2747	293.3	35.54	5
nvsh-15a	New Britain–Vanuatu	159.9736	-9.5993	302.8	46.69	41.38
nvsh-15b	New Britain–Vanuatu	159.8044	-9.8584	302.8	46.69	5
nvsh-16a	New Britain–Vanuatu	160.7343	-10.0574	301	46.05	41
nvsh-16b	New Britain–Vanuatu	160.5712	-10.3246	301	46.05	5
nvsh-17a	New Britain–Vanuatu	161.4562	-10.5241	298.4	40.12	37.22
nvsh-17b	New Britain–Vanuatu	161.2900	-10.8263	298.4	40.12	5
nvsh-18a	New Britain–Vanuatu	162.0467	-10.6823	274.1	40.33	29.03
nvsh-18b	New Britain–Vanuatu	162.0219	-11.0238	274.1	28.72	5
nvsh-19a	New Britain–Vanuatu	162.7818	-10.5645	261.3	34.25	24.14
nvsh-19b	New Britain–Vanuatu	162.8392	-10.9315	261.3	22.51	5
nvsh-20a	New Britain–Vanuatu	163.7222	-10.5014	262.9	50.35	26.3
nvsh-20b	New Britain–Vanuatu	163.7581	-10.7858	262.9	25.22	5
nvsh-21a	New Britain–Vanuatu	164.9445	-10.4183	287.9	40.31	23.3
nvsh-21b	New Britain–Vanuatu	164.8374	-10.7442	287.9	21.47	5
nvsh-22a	New Britain–Vanuatu	166.0261	-11.1069	317.1	42.39	20.78
nvsh-22b	New Britain–Vanuatu	165.7783	-11.3328	317.1	18.4	5
nvsh-23a	New Britain–Vanuatu	166.5179	-12.2260	342.4	47.95	22.43
nvsh-23b	New Britain–Vanuatu	166.2244	-12.3171	342.4	20.4	5
nvsh-24a	New Britain–Vanuatu	166.7236	-13.1065	342.6	47.13	28.52
nvsh-24b	New Britain–Vanuatu	166.4241	-13.1979	342.6	28.06	5
nvsh-25a	New Britain–Vanuatu	166.8914	-14.0785	350.3	54.1	31.16
nvsh-25b	New Britain–Vanuatu	166.6237	-14.1230	350.3	31.55	5
nvsh-26a	New Britain–Vanuatu	166.9200	-15.1450	365.6	50.46	29.05
nvsh-26b	New Britain–Vanuatu	166.6252	-15.1170	365.6	28.75	5
nvsh-27a	New Britain–Vanuatu	167.0053	-15.6308	334.2	44.74	25.46

Continued on next page

Table B.8 – continued

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
nvsz-27b	New Britain–Vanuatu	166.7068	-15.7695	334.2	24.15	5
nvsz-28a	New Britain–Vanuatu	167.4074	-16.3455	327.5	41.53	22.44
nvsz-28b	New Britain–Vanuatu	167.1117	-16.5264	327.5	20.42	5
nvsz-29a	New Britain–Vanuatu	167.9145	-17.2807	341.2	49.1	24.12
nvsz-29b	New Britain–Vanuatu	167.6229	-17.3757	341.2	22.48	5
nvsz-30a	New Britain–Vanuatu	168.2220	-18.2353	348.6	44.19	23.99
nvsz-30b	New Britain–Vanuatu	167.8895	-18.2991	348.6	22.32	5
nvsz-31a	New Britain–Vanuatu	168.5022	-19.0510	345.6	42.2	22.26
nvsz-31b	New Britain–Vanuatu	168.1611	-19.1338	345.6	20.2	5
nvsz-32a	New Britain–Vanuatu	168.8775	-19.6724	331.1	42.03	21.68
nvsz-32b	New Britain–Vanuatu	168.5671	-19.8338	331.1	19.49	5
nvsz-33a	New Britain–Vanuatu	169.3422	-20.4892	332.9	40.25	22.4
nvsz-33b	New Britain–Vanuatu	169.0161	-20.6453	332.9	20.37	5
nvsz-34a	New Britain–Vanuatu	169.8304	-21.2121	329.1	39	22.73
nvsz-34b	New Britain–Vanuatu	169.5086	-21.3911	329.1	20.77	5
nvsz-35a	New Britain–Vanuatu	170.3119	-21.6945	311.9	39	22.13
nvsz-35b	New Britain–Vanuatu	170.0606	-21.9543	311.9	20.03	5
nvsz-36a	New Britain–Vanuatu	170.9487	-22.1585	300.4	39.42	23.5
nvsz-36b	New Britain–Vanuatu	170.7585	-22.4577	300.4	21.71	5
nvsz-37a	New Britain–Vanuatu	171.6335	-22.3087	281.3	30	22.1
nvsz-37b	New Britain–Vanuatu	171.5512	-22.6902	281.3	20	5




Figure B.9: New Zealand–Puysegur Zone unit sources.

Table B.9: Earthquake parameters for New Zealand–Puysegur Subduction Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
nzs-1a	New Zealand–Puysegur	168.0294	-45.4368	41.5	15	17.94
nzs-1b	New Zealand–Puysegur	167.5675	-45.1493	41.5	15	5
nzs-2a	New Zealand–Puysegur	167.3256	-46.0984	37.14	15	17.94
nzs-2b	New Zealand–Puysegur	166.8280	-45.8365	37.14	15	5
nzs-3a	New Zealand–Puysegur	166.4351	-46.7897	39.53	15	17.94
nzs-3b	New Zealand–Puysegur	165.9476	-46.5136	39.53	15	5
nzs-4a	New Zealand–Puysegur	166.0968	-47.2583	15.38	15	17.94
nzs-4b	New Zealand–Puysegur	165.4810	-47.1432	15.38	15	5
nzs-5a	New Zealand–Puysegur	165.7270	-48.0951	13.94	15	17.94
nzs-5b	New Zealand–Puysegur	165.0971	-47.9906	13.94	15	5
nzs-6a	New Zealand–Puysegur	165.3168	-49.0829	22.71	15	17.94
nzs-6b	New Zealand–Puysegur	164.7067	-48.9154	22.71	15	5
nzs-7a	New Zealand–Puysegur	164.8017	-49.9193	23.25	15	17.94
nzs-7b	New Zealand–Puysegur	164.1836	-49.7480	23.25	15	5




Figure B.10: Ryukyu-Kyushu-Nankai Zone unit sources.

Table B.10: Earthquake parameters for Ryukyu–Kyushu–Nankai Subduction
Zone unit sources.

Segment	Description	Longitude(°E)	Latitude(°N)	Strike(°)	Dip(°)	Depth (km)
rnsz-1a	Ryukyu–Nankai	122.6672	23.6696	262	14	11.88
rnsz-1b	Ryukyu–Nankai	122.7332	23.2380	262	10	3.2
rnsz-2a	Ryukyu–Nankai	123.5939	23.7929	259.9	18.11	12.28
rnsz-2b	Ryukyu–Nankai	123.6751	23.3725	259.9	10	3.6
rnsz-3a	Ryukyu–Nankai	124.4604	23.9777	254.6	19.27	14.65
rnsz-3b	Ryukyu–Nankai	124.5830	23.5689	254.6	12.18	4.1
rnsz-4a	Ryukyu–Nankai	125.2720	24.2102	246.8	18	20.38
rnsz-4b	Ryukyu–Nankai	125.4563	23.8177	246.8	16	6.6
rnsz-5a	Ryukyu–Nankai	125.9465	24.5085	233.6	18	20.21
rnsz-5b	Ryukyu–Nankai	126.2241	24.1645	233.6	16	6.43
rnsz-6a	Ryukyu–Nankai	126.6349	25.0402	228.7	17.16	19.55
rnsz-6b	Ryukyu–Nankai	126.9465	24.7176	228.7	15.16	6.47
rnsz-7a	Ryukyu–Nankai	127.2867	25.6343	224	15.85	17.98
rnsz-7b	Ryukyu–Nankai	127.6303	25.3339	224	13.56	6.26
rnsz-8a	Ryukyu–Nankai	128.0725	26.3146	229.7	14.55	14.31
rnsz-8b	Ryukyu–Nankai	128.3854	25.9831	229.7	9.64	5.94
rnsz-9a	Ryukyu–Nankai	128.6642	26.8177	219.2	15.4	12.62
rnsz-9b	Ryukyu–Nankai	129.0391	26.5438	219.2	8	5.66
rnsz-10a	Ryukyu–Nankai	129.2286	27.4879	215.2	17	12.55
rnsz-10b	Ryukyu–Nankai	129.6233	27.2402	215.2	8.16	5.45
rnsz-11a	Ryukyu–Nankai	129.6169	28.0741	201.3	17	12.91
rnsz-11b	Ryukyu–Nankai	130.0698	27.9181	201.3	8.8	5.26
rnsz-12a	Ryukyu–Nankai	130.6175	29.0900	236.7	16.42	13.05
rnsz-12b	Ryukyu–Nankai	130.8873	28.7299	236.7	9.57	4.74
rnsz-13a	Ryukyu–Nankai	130.7223	29.3465	195.2	20.25	15.89
rnsz-13b	Ryukyu–Nankai	131.1884	29.2362	195.2	12.98	4.66
rnsz-14a	Ryukyu–Nankai	131.3467	30.3899	215.1	22.16	19.73
rnsz-14b	Ryukyu–Nankai	131.7402	30.1507	215.1	17.48	4.71
rnsz-15a	Ryukyu–Nankai	131.9149	31.1450	216	15.11	16.12
rnsz-15b	Ryukyu–Nankai	132.3235	30.8899	216	13.46	4.48
rnsz-16a	Ryukyu–Nankai	132.5628	31.9468	220.9	10.81	10.88
rnsz-16b	Ryukyu–Nankai	132.9546	31.6579	220.9	7.19	4.62
rnsz-17a	Ryukyu–Nankai	133.6125	32.6956	239	10.14	12.01
rnsz-17b	Ryukyu–Nankai	133.8823	32.3168	239	8.41	4.7
rnsz-18a	Ryukyu–Nankai	134.6416	33.1488	244.7	10.99	14.21
rnsz-18b	Ryukyu–Nankai	134.8656	32.7502	244.5	10.97	4.7
rnsz-19a	Ryukyu–Nankai	135.6450	33.5008	246.5	14.49	14.72
rnsz-19b	Ryukyu–Nankai	135.8523	33.1021	246.5	11.87	4.44
rnsz-20a	Ryukyu–Nankai	136.5962	33.8506	244.8	15	14.38
rnsz-20b	Ryukyu–Nankai	136.8179	33.4581	244.8	12	3.98
rnsz-21a	Ryukyu–Nankai	137.2252	34.3094	231.9	15	15.4
rnsz-21b	Ryukyu–Nankai	137.5480	33.9680	231.9	12	5
rnsz-22a	Ryukyu–Nankai	137.4161	34.5249	192.3	15	15.4
rnsz-22b	Ryukyu–Nankai	137.9301	34.4327	192.3	12	5

Appendix C. SIFT Testing Report:

Arena Cove, California

Jean Newman

1.0 PURPOSE

Forecast models are tested with synthetic tsunami events covering a range of tsunami source locations. Testing is also done with selected historical tsunami events when available.

The purpose of forecast model testing is three-fold. The first objective is to assure that the results obtained with NOAA's tsunami forecast system, which has been released to the Tsunami Warning Centers for operational use, are identical to those obtained by the researcher during the development of the forecast model. The second objective is to test the forecast model for consistency, accuracy, time efficiency, and quality of results over a range of possible tsunami locations and magnitudes. The third objective is to identify bugs and issues in need of resolution by the researcher who developed the Forecast Model or by the forecast software development team before the next version release to NOAA's two Tsunami Warning Centers.

Local hardware and software applications, and tools familiar to the researcher(s), are used to run the Method of Splitting Tsunamis (MOST) model during the forecast model development. The test results presented in this report lend confidence that the model performs as developed and produces the same results when initiated within the forecast application in an operational setting as those produced by the researcher during the forecast model development. The test results assure those who rely on the Arena Cove tsunami forecast model that consistent results are produced irrespective of system.

2.0 TESTING PROCEDURE

The general procedure for forecast model testing is to run a set of synthetic tsunami scenarios and a selected set of historical tsunami events through the forecast system application and compare the results with those obtained by the researcher during the forecast model development and presented in the Tsunami Forecast Model Report. Specific steps taken to test the model include:

1. Identification of testing scenarios, including the standard set of synthetic events, appropriate historical events, and customized synthetic scenarios that may have been used by the researcher(s) in developing the forecast model.
2. Creation of new events to represent customized synthetic scenarios used by the researcher(s) in developing the forecast model, if any.
3. Submission of test model runs with the forecast system, and export of the results from A, B, and C grids, along with time series.
4. Recording applicable metadata, including the specific version of the forecast system used for testing.
5. Examination of forecast model results from the forecast system for instabilities in both time series and plot results.
6. Comparison of forecast model results obtained through the forecast system with those obtained during the forecast model development.
7. Summarization of results with specific mention of quality, consistency, and time efficiency.
8. Reporting of issues identified to modeler and forecast software development team.
9. Retesting the forecast models in the forecast system when reported issues have been addressed or explained.

Synthetic model runs were tested on a DELL PowerEdge R510 computer equipped with two Xeon E5670 processors at 2.93 Ghz, each with 12 MBytes of cache and 32GB memory. The processors are hex core and support hyperthreading, resulting in the computer performing as a 24 processor core machine. Additionally, the testing computer supports 10 Gigabit Ethernet for fast network connections. This computer configuration is similar or the same as the configurations of the computers installed at the Tsunami Warning Centers so the compute times should only vary slightly.

Results

The Arena Cove forecast model was tested with NOAA's tsunami forecast system version 3.2. The propagation database used during development is consistent with this version.

The Arena Cove, California forecast model was tested with five synthetic scenarios and one historical tsunami event. Test results from the forecast system and comparisons with the results obtained during the forecast model development are shown numerically in Table C1 and graphically in Figures C1 to C4. The results show that the forecast model is stable and robust, with consistent and high quality results across geographically distributed tsunami sources and mega-event tsunami magnitudes. The model run time (wall clock time) was under 19 minutes for 8 hours of simulation time, and under 10 minutes for 4 hours. This run time is within the 10 minute run time for 4 hours of simulation time and satisfies time efficiency requirements.

Five synthetic events (there were no timeseries plots in the report for 2 of the cases) were run on the Arena Cove forecast model. The modeled scenarios were stable for all cases tested, with no instabilities or ringing. Results show that the largest modeled height was 295.03 cm and originated in the Aleutian-Alaska-Cascadia (ACSZ 56-65) source. Amplitudes greater than 100 cm were recorded for 4 out of 5 test sources. The smallest signal of 47.67 cm was recorded at the far field Central and South American (CSSZ 89-98) source. Direct comparisons, of output from the forecast tool with results of both the historical event (Tohoku, Honshu in report, 2011) and available development synthetic events, demonstrated that the wave pattern were similar in shape, pattern and amplitude.

Table 1. Table of maximum and minimum amplitudes (cm) at the Arena Cove, California warning point for synthetic and historical events tested using SIFT 3.2 and obtained during development.

Scenario Name	Source Zone	Tsunami Source	α [m]	SIFT Max (cm)	Development Max (cm)	SIFT Min (cm)	Development Min (cm)
Mega-tsunami Scenarios							
KISZ 1-10	Kamchatka-Yap-Mariana-Izu-Bonin	A1-A10, B1-B10	25	250.055	250.055	-368.491	-315.426
KISZ 22-31	Kamchatka-Yap-Mariana-Izu-Bonin	A22-A31, B22-B31	25	266.764	266.764	-198.325	-198.325
ASCZ 56-65	Aleutian-Alaska-Cascadia	A56-A65, B56-B65	25	295.027	292.952	-226.995	-231.452
CSSZ 89-98	Central and South America	A89-A98, B89-B98	25	47.673	47.673	-47.978	-47.978
NTSZ 30-39	New Zealand-Kermadec-Tonga	A30-A39, B30-B39	25	175.210	175.21	-179.082	-179.082
Historical Events							
Tohoku 2011	Kamchatka-Yap-Mariana-Izu-Bonin	4.66 b24 + 12.23 b25+26.31 a26+21.27 b26+22.75 a27 +4.98 b27		147.809	147.957	-133.476	-133.518




Figure C1 Response of the Arena Cove forecast model to synthetic scenario ACSZ 56-65. (a,b, and c) Maximum sea surface elevation for A-, B- and C-grids. (d) Sea surface elevation time series at the C-grid warning point.




Figure C2 Response of the Arena Cove forecast model to synthetic scenario CSSZ 89-98. (a,b, and c) Maximum sea surface elevation for A-, B- and C-grids. (d) Sea surface elevation time series at the C-grid warning point.




Figure C3 Response of the Arena Cove forecast model to synthetic scenario NTSZ 30-39. (a,b, and c) Maximum sea surface elevation for A-, B- and C-grids. (d) Sea surface elevation time series at the C-grid warning point.




Figure C4 Response of the Arena Cove forecast model to synthetic scenario NTSZ 30-39. (a, b, and c) Maximum sea surface elevation for A-, B- and C-grids. (d) Sea surface elevation time series at the C-grid warning point.